

## References

For a comprehensive bibliography (6062 references!) about hypotrichs and euplotids, see Berger (2006a).

- Acosta-Mercado D., Lynn D.H. (2003): The edaphic quantitative protargol stain: a sampling protocol for assessing soil ciliate abundance and diversity. – *J. Microbiol. Methods* **53**: 365–375.
- Adam H., Czihak G. (1964): Arbeitsmethoden der makroskopischen und mikroskopischen Anatomie. Ein Laboratoriumshandbuch für Biologen, Mediziner und technische Hilfskräfte. Fischer Verlag, Stuttgart. I–XIV, 583 pp.
- Adl S.M., Simpson A.G.B., Farmer M.A., Andersen R.A., Anderson O.R., Barta J.R., Bowser S.S., Bruggenolte G., Fensome R.A., Fredericq S., James T.Y., Karpov S., Kugrens P., Krug J., Lane C.E., Lewis L.A., Lodge J., Lynn D.H., Mann D.G., Mccourt R.M., Mendoza L., Moestrup O., Mozley-Standridge S.E., Nerad T.A., Shearer C.A., Smirnov A.V., Spiegel F.W., Taylor M.F.J.R. (2005): The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. – *J. Euk. Microbiol.* **52**: 399–451.
- Aescht E. (2001): Catalogue of the generic names of ciliates (Protozoa, Ciliophora). – *Denisia* **1**: 1–350.
- Aescht E. (2003): Typen-Liste der Sammlung “Wirbellose Tiere” (ohne Insekten) am Biologiezentrum Linz. – *Beitr. Naturk. Oberösterreichs* **12**: 377–406.
- Aescht E., Foissner W. (1993): Effects of organically enriched magnesite fertilizers on the soil ciliates of a spruce forest. – *Pedobiologia* **37**: 321–335.
- Agamaliyev F.G. (1967): Faune des ciliés mésopsammiques de la côte ouest de la mer Caspienne. – *Cah. Biol. mar.* **8**: 359–402.
- Agamaliyev F.G. (1967a): Ecology of psammophilous ciliates. – *Dokl. Akad. Nauk SSSR* **176**: 1425–1427 (in Russian).
- Agamaliyev F.G. (1970): Vertical distribution of psammophilous ciliates in the Caspian Sea. – *Zool. Zh.* **49**: 1277–1284 (in Russian with English summary).
- Agamaliyev F.G. (1971): Complements to the fauna of psammophilic ciliates of the Western coast of the Caspian Sea. – *Acta Protozool.* **8**: 379–404, Plates I, II (in Russian with English summary).
- Agamaliyev F.G. (1972): Ciliates from microbenthos of the islands of Apseronskij and Bakinskij archipelagos of the Caspian Sea. – *Acta Protozool.* **10**: 1–27 (in Russian with English summary).
- Agamaliyev F.G. (1973): Ciliates of microbenthos in the Krasnovodsk Bay of the Caspian Sea. – *Zool. Zh.* **52**: 1597–1601 (in Russian with English summary).
- Agamaliyev F.G. (1974): Ciliates of the solid surface overgrowth of the Caspian Sea. – *Acta Protozool.* **13**: 53–83, Plates I–IV (in Russian with English summary).
- Agamaliyev F.G. (1974a): Ciliates of the Turkmen Bay (Caspian Sea). – *Zool. Zh.* **53**: 19–24 (in Russian with English summary).
- Agamaliyev F.G. (1983): Infuzorii Kaspijskogo morja. Sistematika, ekologija, zoogeografija (Ciliates of the Caspian Sea. Systematics, ecology, zoogeography). Nauka, Leningrad (in Russian). 232 pp, Plates I–XX.
- Agamaliyev F.G. (1986): Ciliates of the low-salinity lagoons of the Caspian Sea. – *Arch. Protistenk.* **131**: 201–214.
- Agamaliyev F.G. (1990): The fauna and distribution of ciliates in supralittoral zone of the Apsheron Coast of the Caspian Sea. – *Protozoology* **13**: 58–63 (in Russian with English summary).
- Agamaliyev F.G. (1974): Benthic infusoria in the western Caspian Sea. – *Hydrobiol. J.* **10**: 18–23.
- Agamaliyev F.G. (1976): Planctonic and periphytonic ciliophora in the eastern part of the Middle Caspian. – *Izv. Akad. Nauk azerb. SSR (Biologičeskie)* year **1976**: 90–94 (in Russian).
- Agamaliyev F.G., Aliyev A.R. (1983): Benthic infusoria from the Divichinskiy estuary of the Caspian Sea. – *Hydrobiol. J.* **18** (years 1982/1983): 20–24.
- Agatha S. (2003): Morphology and ontogenesis of *Novistrombidium apsheronicum* nov. comb. and *Strombidium arenicola* (Protozoa, Ciliophora): a comparative light microscopical and SEM study. – *Europ. J. Protistol.* **39**: 245–266.

- Agatha S. (2004): A cladistic approach for the classification of oligotrichid ciliates (Ciliophora: Spirotricha). – *Acta Protozool.* **43**: 201–217.
- Agatha S., Riedl-Lorje J.C. (2006): Redescription of *Tintinnopsis cylindrica* Daday, 1887 (Ciliophora: Spirotricha) and unification of tintinnid terminology. – *Acta Protozool.* **45**: 137–151.
- Agatha S., Strüder-Kypke M.C. (2007): Phylogeny of the order Choreotrichida (Ciliophora, Spirotricha, Oligotrichea) as inferred from morphology, ultrastructure, ontogenesis, and SSrRNA gene sequences. – *Europ. J. Protistol.* **43**: 37–63.
- Agatha S., Wilbert N., Spindler M., Elbrächter M. (1990): Euplotide ciliates in sea ice of the Weddell Sea (Antarctica). – *Acta Protozool.* **29**: 221–228.
- Agatha S., Strüder-Kypke M.C., Beran A. (2004): Morphologic and genetic variability in the marine planktonic ciliate *Laboea strobila* Lohmann, 1908 (Ciliophora, Oligotrichia), with notes on its ontogenesis. – *J. Euk. Microbiol.* **51**: 267–281.
- Agatha S., Strüder-Kypke M.C., Beran A., Lynn D.H. (2005): *Pelagostrobilidium neptuni* (Montagnes and Taylor, 1994) and *Strombidium biarmatum* nov. spec. (Ciliophora, Oligotrichea): phylogenetic position inferred from morphology, ontogenesis, and gene sequence data. – *Europ. J. Protistol.* **41**: 65–83.
- Aladro Lubel M.A. (1985): Algunos ciliados intersticiales de Isla de Enmedio, Veracruz, México. – *An. Inst. Biol. Univ. Méx., Ser. Zoología* (1) **55** (year 1984): 1–59, Lamina 1–17.
- Aladro-Lubel M.A., Martín-Murillo M.E., Mayén-Estrada R., Hernández-Anaya M., Sánchez-Calderón G. (1986): Cuarenta y nueve especies de ciliados intersticiales identificados en Boca del Río, Veracruz, México. – *Revta lat.-am. Microbiol.* **28**: 231–242.
- Aladro-Lubel M.A., Martínez-Murillo M.E., Mayén-Estrada R. (1988): Lista de los ciliados bentónicos salobres y marinos registrados en México. – *An. Inst. Biol. Univ. Méx., Ser. Zoología* (1) **58** (year 1987): 403–447.
- Aladro Lubel M.A., Martínez Murillo M.E., Mayén Estrada R. (1990): Manual de ciliados psamófilos marinos y salobres de México. Cuadernos del Instituto de Biología 9. Universidad Nacional Autónoma de México. 174 pp.
- Alekperov I.K. (2005): Atlas free-living infusoria. Baku (in Russian). 310 pp.
- Alekperov I.K., Asadullayeva E.S. (1999): New, little-known and characteristic ciliate species from the Apsheron coast of the Caspian Sea. – *Turkish J. Zool.* **23**: 215–225.
- Alekperov I.H., Musayev M.A. (1988): New and rare free-living infusoria from fresh waters and soil of the Apsheron peninsula. – *Zool. Zh.* **67**: 1904–1909 (in Russian with English summary).
- AL-Rasheid K.A.S. (1999a): Free-living marine interstitial hypotrichid ciliates from Jubail marine wildlife sanctuary in the Arabian Gulf. – *Agricultural Sciences* **4**: 53–62.
- AL-Rasheid K.A.S. (1999b): A review of marine and brackish water interstitial ciliates from the Arabian Gulf, its offshore islands and Al-Hassa Oasis with notes on their ecological status and recovery after the 1991 Gulf War oil spill. – *Arab. Gulf J. Scient. Res.* **17**: 336–368.
- AL-Rasheid K.A.S. (2001): New records of interstitial ciliates (Protozoa Ciliophora) from the Saudi coasts of the Red Sea. – *Tropical Zoology* **14**: 133–156.
- Ammermann D. (1990): The contribution of hypotrichous ciliates to our understanding of molecular biology and evolution of ciliates. – *Zool. Sci. Suppl.* **7**: 13–22.
- Andelová K., Tirjaková E. (2000): Nálevníky (Ciliophora) podložia machových zrástov na územi Bratislavy (Ciliates (Ciliophora) of the underlying moss-covered stratum on the region of Bratislava city. – *Folia faunistica Slovaca* **5**: 27–36 (in Slovakian with English summary).
- AOÖLR (Amt der Oberösterreichischen Landesregierung) (1995): Trattnach und Innbach, Untersuchungen zur Gewässergüte. Stand 1992–1994. – *Gewässerschutz Bericht* **11**: 1–137.
- Ashhurst D.E. (1959): The connective tissue sheath of the locust nervous system: a histochemical study. – *Quart. J. microsc. Sci.* **100**: 401–412.
- Ax P. (1984): Das phylogenetische System. Systematisierung der lebenden Natur aufgrund ihrer Phylogenese. Gustav Fischer Verlag, Stuttgart New York. 349 pp.
- Ax P. (1995): Das System der Metazoa I. Ein Lehrbuch der phylogenetischen Systematik. G. Fischer Verlag, Stuttgart Jena New York. 226 pp.
- Ax P., Ax R. (1960): Experimentelle Untersuchungen über die Salzgehaltstoleranz von Ciliaten aus dem Brackwasser und Süßwasser. – *Biol. Zbl.* **79**: 7–31.

- Azovskii A.I., Mazei Y.A. (2003): Ciliates of coarse ground on the northeastern Black Sea coast. – Zool. Zh. **82**: 899–912 (in Russian with English summary).
- Azovsky A.I. (1996): New data on the fauna of benthic infusorians (Ciliata, Protozoa) of the Barents Sea. – Zool. Zh. **75**: 3–8 (in Russian with English summary).
- Azovsky A.I., Mazei Y.A. (2003): A conspectus of the Black Sea fauna of benthic ciliates. – Protistology **3**: 72–91.
- Azovsky A.I., Mazei Y.A. (2005): Distribution and community structure of benthic ciliates in the North-Eastern part of the Black Sea. – Protistology **4**: 83–90.
- Azovsky A.I., Mazei Y.A. (2007): Structure of subtidal and intertidal communities of psammophilous ciliates of the Pechora Sea. – Oceanology **47**: 60–67.
- Azovsky A.I., Burkovsky I.V., Kashunin A.K., Akseonov D.E. (1996): The White Sea littoral micro-benthic communities as an indicator of environmental conditions. – Russian Journal aquatic Ecology **5**: 27–37.
- Babko R.V., Kovalchuk A.A.** (1992): Free-living infusoria of the Sumsoi region. In: Ministry of Education & Sumsoi, Pedagogical Institute Makarenko of the Town (eds.): Problems of protection and conservation of the natural resources of the Sumsoi region, pp. 113–125. Sumsoi (in Ukrainian).
- Băcescu M., Dumitrescu E., Gomoiu M.T., Petran A. (1967): Éléments pour la caractérisation de la zone sédimentaire médio- littorale de la Mer Noire. – Trav. Mus. Hist. nat. Gr. Antipa **7**: 1–14.
- Bamforth S.S. (1968): Forest soil protozoa of the Florida Parishes of Louisiana. – Proc. La Acad. Sci. **31**: 5–15.
- Bamforth S.S. (2004): Water film fauna of microbiotic crusts of a warm desert. – J. Arid Environments **56**: 413–423.
- Bardele C.F. (1981): Functional and phylogenetic aspects of the ciliary membrane: a comparative freeze-fracture study. – BioSystems **14**: 403–421.
- Berg J.M., Tymoczko J.L., Stryer L. (2003): Biochemie. Fifth edition. Spektrum Akademischer Verlag, Heidelberg, Berlin. i–xxxv, 1153 pp.
- Berger H. (1985): Taxonomie und Ökologie der Ciliaten und Testaceen (Protozoa) von Almweiden und Schipisten im Gasteinertal (Salzburg). Dissertation, Universität Salzburg. 158 pp.
- Berger H. (1999): Monograph of the Oxytrichidae (Ciliophora, Hypotrichia). – Monographiae biol. **78**: i–xii, 1–1080.
- Berger H. (2001): Catalogue of ciliate names 1. Hypotrichs. Helmut Berger, Salzburg. i–viii, 206 pp.
- Berger H. (2003): Redefinition of *Holosticha* Wrzesniowski, 1877 (Ciliophora, Hypotricha). – Europ. J. Protistol. **39**: 373–379.
- Berger H. (2004): *Amphisiella annulata* (Kahl, 1928) Borror, 1972 (Ciliophora: Hypotricha): morphology, notes on morphogenesis, review of literature, and neotypification. – Acta Protozool. **43**: 1–14.
- Berger H. (2004a): *Uroleptopsis* Kahl, 1932 (Ciliophora: Hypotricha): morphology and cell division of type species, redefinition, and phylogenetic relationships. – Acta Protozool. **43**: 99–121.
- Berger H. (2005a): Monograph of the hypotrichs (Ciliophora). Part 3. - A biodiversity study. – Jahrestagung der Deutschen Gesellschaft für Protozoologie **24** (Kaiserslautern): Abstract.
- Berger H. (2005b): Revision of the Amphisiellidae Jankowski, 1979 (Ciliophora, Hypotricha). – Int. Congr. Protozool. **12**: 106, Abstract S4.1-25.
- Berger H. (2006): Monograph of the Urostyloidea (Ciliophora, Hypotricha). – Monographiae biol. **85**: i–xvi, 1–1304.
- Berger H. (2006a): Bibliography of Hypotrichs and Euplotids (Ciliophora). Verlag Helmut Berger, Salzburg. i–viii, 425 pp.
- Berger H. (2007): The ground-pattern of the Hypotricha (Spirotricha, Ciliophora). – Protistology **5**: 14, Abstract.
- Berger H. (2008): How did the last common ancestor of the hypotrichs (Ciliophora, Spirotricha) look like? – Jahrestagung der Deutschen Gesellschaft für Protozoologie **27** (Warnemünde, Rostock): Abstract on p. 30 of abstract booklet.
- Berger H., Foissner W. (1987): Morphology and biometry of some soil hypotrichs (Protozoa: Ciliophora). – Zool. Jb. Syst. **114**: 193–239.

- Berger H., Foissner W. (1987a): Guide-lines for the alpha-taxonomy of hypotrichous ciliates. – Society of Protozoologists **1986 Abstracts**: Abstract 136.
- Berger H., Foissner W. (1988): Revision of *Lamostyla* Buitkamp, 1977 and description of *Territricha* nov. gen. (Ciliophora: Hypotrichida). – Zool. Anz. **220**: 113–134.
- Berger H., Foissner W. (1989): Morphology and biometry of some soil hypotrichs (Protozoa, Ciliophora) from Europe and Japan. – Bull. Br. Mus. nat. Hist. (Zool.) **55**: 19–46.
- Berger H., Foissner W. (1989a): Morphology and morphogenesis of *Parakahliella haideri* nov. spec. (Ciliophora, Hypotrichida). – Bull. Br. Mus. nat. Hist. (Zool.) **55**: 11–17.
- Berger H., Foissner W. (1992): Polyhymenophora Jankowski, 1967: a new name for the spirotrichs of Bütschli? – J. Protozool. Suppl. **39**: 13A, Abstract 77.
- Berger H., Foissner W. (1997): Cladistic relationships and generic characterization of oxytrichid hypotrichs (Protozoa, Ciliophora). – Arch. Protistenk. **148**: 125–155.
- Berger H. Foissner W. (2003): Illustrated guide and ecological notes to ciliate indicator species (Protozoa, Ciliophora) in running waters, lakes, and sewage plants. In: Steinberg, Wilken, Calmano, Klapper (eds.): Handbuch angewandte Limnologie, Vol. **17**. Erg.Lfg, pp. 1–160. ecomed, Landsberg am Lech.
- Berger H., Foissner W., Adam H. (1983): Morphology and morphogenesis of *Fuscheria terricola* n. sp. and *Spathidium muscorum* (Ciliophora: Kinetofragminophora). – J. Protozool. **30**: 529–535.
- Berger H., Foissner W., Adam H. (1984): Taxonomie, Biometrie und Morphogenese einiger terricoler Ciliaten (Protozoa: Ciliophora). – Zool. Jb. Syst. **111**: 339–367.
- Berger H., Foissner W., Adam H. (1985): Protozoologische Untersuchungen an Almböden im Gasteiner Tal (Zentralalpen, Österreich). IV. Experimentelle Studien zur Wirkung der Bodenverdichtung auf die Struktur der Testaceen- und Ciliatentaxozönose. – Veröff. Österr. MaB-Programms **9**: 97–112.
- Berger H., Foissner W., Adam H. (1985a): Morphological variation and comparative analysis of morphogenesis in *Parakahliella macrostoma* (Foissner, 1982) nov. gen. and *Histiculus muscorum* (Kahl, 1932), (Ciliophora, Hypotrichida). – Protistologica **21**: 295–311.
- Berger H., Foissner W., Adam H. (1986): Field experiments on the effects of fertilizers and lime on the soil microfauna of an alpine pasture. – Pedobiologia **29**: 261–272.
- Berger H., Foissner W. Kohmann F. (1997): Bestimmung und Ökologie der Mikrosaprobien nach DIN 38410. Gustav Fischer Verlag, Stuttgart, Jena, Lübeck, Ulm. I–X & 291 pp.
- Bernhard D., Stechmann A., Foissner W., Ammermann D., Hehn M., Schlegel M. (2001): Phylogenetic relationships within the class Spirotrichea (Ciliophora) inferred from small subunit rRNA gene sequences. – Molecular Phylogenetics and Evolution **21**: 86–92.
- Bick H., Buitkamp U. (1976): Ciliated protozoa from Canadian grassland soils. – Trans. Am. microsc. Soc. **95**: 490–491.
- Biczók F. (1956): Contributions to the protozoa of the rhizosphere of wheat. – Acta zool. hung. **2**: 115–147.
- Biernacka I. (1962): Die Protozoenfauna in der Danziger Bucht I. Die Protozoen in einigen Biotopen der Seeküste. – Polskie Archwm Hydrobiol. **10**: 39–109.
- Biernacka I. (1963): Die Protozoenfauna in Danziger Bucht II. Die Charakteristik der Protozoen in untersuchten Biotopen der Seeküste. – Polskie Archwm Hydrobiol. **11**: 17–75.
- Blackburn E.H. (1990): Telomeres: structure and synthesis. – J. biol. Chem. **265**: 5919–5921.
- Blatterer H., Foissner W. (1988): Beitrag zur terricolen Ciliatenfauna (Protozoa: Ciliophora) Australiens. – Stapfia **17**: 1–84.
- Blatterer H., Foissner W. (2003): Morphological and ontogenetic comparison of two populations of *Parentocirrus hortualis* Voss 1997 (Ciliophora, Hypotrichida). – Linzer biol. Beitr. **35**: 831–854.
- Bock K.J. (1952): Zur Ökologie der Ciliaten des marinen Sandgrundes der Kieler Bucht I. – Kieler Meeressforsch. **9**: 77–89.
- Bock K.J. (1953): Zur Oekologie der Ciliaten des marinen Sandgrundes der Kieler Bucht II. – Kieler Meeressforsch. **9**: 252–256.
- Bock K.J. (1960): Biologische Untersuchungen, insbesondere der Ciliatenfauna, in der durch Abwässer belasteten Schlei (westliche Ostsee). – Kieler Meeressforsch. **16**: 57–68.

- Bocquet C. (1971): Espèces nouvelles décrites de la région de Roscoff entre 1945 et 1970. – Cah. Biol. mar. **12**: 381–404.
- Borror A.C. (1962): Ciliate protozoa of the Gulf of Mexico. – Bull. mar. Sci. Gulf Caribb. **12**: 333–349.
- Borror A.C. (1963): Morphology and ecology of the benthic ciliated protozoa of Alligator Harbor, Florida. – Arch. Protistenk. **106**: 465–534.
- Borror A.C. (1963a): Morphology and ecology of some uncommon ciliates from Alligator Harbor, Florida. – Trans. Am. microsc. Soc. **82**: 125–131.
- Borror A.C. (1972): Revision of the order Hypotrichida (Ciliophora, Protozoa). – J. Protozool. **19**: 1–23.
- Borror A. (1972a): Tidal marsh ciliates (Protozoa): morphology, ecology, systematics. – Acta Protozool. **10**: 29–71, Plate I.
- Borror A.C. (1973): Marine flora and fauna of the Northeastern United States. Protozoa: Ciliophora. – NOAA Technical Report NMFS CIRC-378: i–iv, 1–62.
- Borror A.C. (1979): Redefinition of the Urostylidae (Ciliophora, Hypotrichida) on the basis of morphogenetic characters. – J. Protozool. **26**: 544–550.
- Borror A.C., Evans F.R. (1979): *Cladotricha* and phylogeny in the suborder Stichotrichina (Ciliophora, Hypotrichida). – J. Protozool. **26**: 51–55.
- Borror A.C., Hill B.F. (1995): The order Euplotida (Ciliophora): Taxonomy, with division of *Euplotes* into several genera. – J. Euk. Microbiol. **42**: 457–466.
- Borror A.C., Wicklow B.J. (1982): Non-homology of median rows in hypotrichs with only three longitudinal rows of cirri. – J. Protozool. Suppl. **29**: 285A, Abstract.
- Borror A.C., Wicklow B.J. (1983): The suborder Urostylina Jankowski (Ciliophora, Hypotrichida): morphology, systematics and identification of species. – Acta Protozool. **22**: 97–126, Plate I.
- Brugerolle G., Lee J.J. (2002): Phylum Parabasalia. In: Lee J.J., Leedale G.F., Bradbury P. (eds.): An illustrated guide to the Protozoa. Second edition. Organisms traditionally referred to as Protozoa, or newly discovered groups. Volume I, pp. 1196–1250. Society of Protozoologists, Lawrence.
- Buitkamp U. (1977): Die Ciliatenfauna der Savanne von Lamto (Elfenbeinküste). – Acta Protozool. **16**: 249–276.
- Buitkamp U. (1977a): Über die Ciliatenfauna zweier mitteleuropäischer Bodenstandorte (Protozoa; Ciliata). – Decheniana **130**: 114–126.
- Buitkamp U. (1979): Vergleichende Untersuchungen zur Temperaturadaptation von Bodenciliaten aus klimatisch verschiedenen Regionen. – Pedobiologia **19**: 221–236.
- Buitkamp U., Wilbert N. (1974): Morphologie und Taxonomie einiger Ciliaten eines kanadischen Präriebodens. – Acta Protozool. **13**: 201–210.
- Bullington W.E. (1925): A study of spiral movement in the ciliate infusoria. – Arch. Protistenk. **50**: 219–274.
- Burkovskii I.V., Azovskii A.I. (1985a): A reverse succession in a community of marine psammophilous ciliates in the laboratory experiment. – Zool. Zh. **64**: 615–618 (in Russian with English summary).
- Burkovskii I.V., Azovskii A.I. (1985b): On correlation between the species numbers and stability in a community of marine psammophilous ciliates. – Zool. Zh. **64**: 813–819 (in Russian with English summary).
- Burkovsky I.V. (1968): Quantitative data on the vertical distribution of psammophylic infusoria in the Velikaya Salma (Kandalaksha Bay, The White Sea). – Zool. Zh. **47**: 1407–1410 (in Russian with English summary).
- Burkovsky I.V. (1970): Psammophilous infusoria of the Kandalakshskiy Bay (White Sea). Dissertation, Moscow University Publisher. 21 pp. (in Russian).
- Burkovsky I.V. (1970a): Ciliates of the mesopsammon of the littoral and sublittoral areas in the Bay of Kandalaksha. – Trudy Belomorskoi Biologicheskoi Stantsii. M. G. U. **3**: 51–59 (in Russian).
- Burkovsky I.V. (1970b): Ciliates of the sand littoral and sublittoral of Kandalaksha Gulf (White Sea) and the analysis on the fauna of benthic ciliates of other seas. – Acta Protozool. **8**: 183–201 (in Russian with English summary).
- Burkovsky I.V. (1971a): Ciliates of the rocky supralittoral zone in the White Sea. – Zool. Zh. **50**: 1569–1572 (in Russian with English summary).

- Burkovsky I.V. (1971b): A comparative study of the ecology of free-living ciliates in the Rugozersky inlet (Kandalaksha Bay, White Sea). – *Zool. Zh.* **50**: 1773–1779 (in Russian with English summary).
- Burkovsky I.V. (1971c): Ecology of psammophilous ciliates in the White Sea. – *Zool. Zh.* **50**: 1285–1302 (in Russian with English summary).
- Burkovsky I.V. (1976): Ciliates of the freshened sites of the White Sea. – *Zool. Zh.* **55**: 287–289 (in Russian with English summary).
- Burkovsky I.V. (1978): Structure, dynamics and production of a community of marine psammophilous ciliates. – *Zool. Zh.* **57**: 325–337 (in Russian with English summary).
- Burkovsky I.V. (1984): *Ekologiya svobodnozivushchikh infuzorij* (Ecology of free-living ciliates). Moscow University Press, Moscow (in Russian). Translation of title from the book review of Seravin L.N. & Raikov I.B., 1985, *Arch. Protistenk.* **129**: 293–294). 208 pp.
- Burkovsky I.V. (1987): The partitioning of ecological resources and the species interrelations in the community of marine psammophilous infusorians. – *Zool. Zh.* **66**: 645–654 (in Russian with English summary).
- Burkovsky I.V., Epstein V.S., Moliboga N.N. (1980): Food specialization and trophic structure of a community of marine psammophilous infusoria. – *Zool. Zh.* **59**: 325–334 (in Russian with English summary).
- Butschinsky P. (1897): Die Protozoen-Fauna der Salzsee-Limane bei Odessa. – *Zool. Anz.* **20**: 194–197.
- Bütschli O. (1889): Protozoa. III. Abtheilung: Infusoria und System der Radiolaria. In: Bronn H.G. (ed.): *Klassen und Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in Wort und Bild, Erster Band*, pp. 1585–2035, Tafeln LVI–LXXIX. Winter, Leipzig.
- Calkins G.N. (1909): *Protozoölogy*. Lea & Febiger, New York, Philadelphia. 349 pp.
- Calvo P., Serrano S., Fernández-Galiano D., Arregui L., Campos I. (1992): Ultrastructural aspects of the mature resting cyst and encystment process in *Phacodinium metchnikoffi*. – *Europ. J. Protistol.* **28**: 334, Abstract 20.
- Calvo P., Fernandez-Aliseda M.C., Garrido J., Torres A. (2003): Ultrastructure, encystment and cyst wall composition of the resting cyst of the peritrich ciliate *Opisthonecta henneguyi*. – *J. Eukaryot. Microbiol.* **50**: 49–56.
- Carey P.G. (1992): *Marine interstitial ciliates. An illustrated key*. Chapman & Hall, London New York Tokyo Melbourne Madras. 351 pp.
- Carey P.G., Maeda M. (1985): Horizontal distribution of psammophilic ciliates in fine sediments of the Chichester harbour area. – *J. nat. Hist. (London)* **19**: 555–574.
- [CBE] Council of Biology Editors (1996): *Scientific Style and format. The CBE manual for authors, editors, and publishers*. 6th edition. Cambridge University Press, Cambridge. 825 pp.
- Chang W.-J., Bryson P., Landweber L. (2003): Searching for the origin of ciliate scrambled genes: hints from several DNA polymerase alpha genes. – International Conference on the Molecular Biology and Biotechnology of Ciliates and Anaerobic Protozoa in connection with 22nd Annual meeting of the German Society for Protozoology, Nijmegen (March 4–8, 2003).
- Chaouite J., Groliere C.A., Pepin D. (1990): Etude du peuplement infusorien des eaux minérales et thermominérales en Auvergne. – *J. Protozool. Suppl.* **37**: 31A, Abstract 149.
- Chen Z., Song W. (2002): Phylogenetic positions of *Aspidisca steini* and *Euplotes vannus* within the order Euplotida (Hypotrichia: Ciliophora) inferred from complete small subunit ribosomal RNA gene sequences. – *Acta Protozool.* **41**: 1–9.
- Chrenková Z., Tirjaková E. (2000): Nálevníky (Ciliophora) suchých machov Bratislavy (Ciliates [Ciliophora] from dry mosses of Bratislava [Slovakia]). – *Folia faunistica Slovaca* **5**: 37–48 (in Slovakian with English abstract).
- Claparède É., Lachmann J. (1858): Études sur les infusoires et les rhizopodes. – *Mém. Inst. natn. génev.* **5** (year 1857): 1–260, Planches 1–13.
- Cohn F. (1866): Neue Infusorien im Seeaquarium. – *Z. wiss. Zool.* **16**: 253–302, Tafeln XIV, XV.
- Cole T.C.H. (2000): *Wörterbuch der Tiernamen*. Spektrum Akademischer Verlag, Heidelberg. 970 pp.

- Corliss J.O. (1960): The problem of homonyms among generic names of ciliated protozoa, with proposal of several new names. – *J. Protozool.* **7**: 269–278.
- Corliss J.O. (1961): The ciliated protozoa: characterization, classification, and guide to the literature. In: Kerker G.A. (ed.): International series of monographs on pure and applied biology, Zoology division Vol. 7, Pergamon Press, Oxford, London, New York, Paris. 310 pp.
- Corliss J.O. (1977): Annotated assignment of families and genera to the orders and classes currently comprising the Corlissian scheme of higher classification for the phylum Ciliophora. – *Trans. Am. microsc. Soc.* **96**: 104–140.
- Corliss J.O. (1979): The ciliated protozoa. Characterization, classification and guide to the literature. Pergamon Press, Oxford, New York, Toronto, Sydney, Paris, Frankfurt. I–XVI, 455 pp.
- Corliss J.O., Lom J. (1985): An annotated glossary of protozoological terms. In: Lee J.J., Hutner S.H., Bovee E.C. (eds.). An illustrated guide to the protozoa, pp. 576–602. Society of Protozoologists, Lawrence.
- Corliss J.O., Lom J. (2002): An annotated glossary of protozoological terms. In: Lee J.J., Leedale G.F., Bradbury P. (eds.). An illustrated guide to the protozoa, second edition, pp. 1346–1385. Society of Protozoologists, Lawrence.
- Cowling A.J. (1994): Protozoan distribution and adaptation. In: Darbyshire J.F. (ed.): Soil protozoa, pp. 5–42. CAB International, Wallingford. I–X, 209 pp.
- Croft K.E., Dalby A.B., Hogan D.J., Orr K.E., Hewitt E.A., Africa R.J., DuBois M.L., Prescott D.M. (2003): Macronuclear molecules encoding actins in spirotrichs. – *J. molec. Evolut.* **56**: 341–350.
- Cunha A.M.da (1916): Contribuição para o conhecimento da fauna de protozoários do Brasil IV. – *Mems Inst. Oswaldo Cruz* **8**: 66–72, estampa 3.
- Curds C.R. (1975): A guide to the species of the genus *Euplotes* (Hypotrichida, Ciliata). – *Bull. Br. Mus. nat. Hist. (Zool.)* **28**: 1–61.
- Curds C.R. (1975a): Protozoa. In: Curds C.R., Hawkes H.A. (eds.): Ecological aspects of used-water treatment. Volume 1 – The organisms and their ecology, pp. 203–268. Academic Press, London, New York, San Francisco.
- Curds C.R., Wu I.C.H. (1983): A review of the Euplotidae (Hypotrichida, Ciliophora). – *Bull. Br. Mus. nat. Hist. (Zool.)* **44**: 191–247.
- Curds C.R., Gates M.A., Roberts D.M. (1983): British and other freshwater ciliated protozoa Part II Ciliophora: Oligohymenophora and Polyhymenophora. Keys and notes for the identification of the free-living genera. In: Kermack D.M., Barnes R.S.K. (eds.): Synopses of the British fauna, No. 23, pp. 1–474. Cambridge University Press, Cambridge, London, New York, New Rochelle, Melbourne, Sydney.
- Czapik A. (1952): Untersuchungen über die Infusorien und Rotatorien des Küstengrundwassers und des Sandbodens der Stalin- Bucht. – *Trud. morsk. biol. Sta. Stalin* **17**: 61–66.
- Czapik A. (1961): Fauna des Kläranlagenmodells in Kraków. – *Acta hydrobiol., Kraków* **3**: 63–67.
- Czapik A. (1962): Mikrofauna słonawego jeziora Ptasi Raj (La microfaune du lac saumâtre Ptasi Raj). – *Polskie Archiwum Hydrobiol.* **10**: 371–378 (in Polish with Russian and French summary).
- Czapik A. (1981): La morphogénèse chez le cilié *Diophrys oligothrix* Borror. – *Acta Protozool.* **20**: 367–372, Planches I, II.
- Czapik A., Fyda J. (1992): Contribution à la connaissance des ciliés psammophiles de la Baltique. – *Acta Protozool.* **31**: 109–114.
- Czapik A., Jordan A. (1976): Les ciliés psammophiles de la mer Baltique aux environs de Gdansk. – *Acta Protozool.* **15**: 423–445, Planches I–IV.
- Dagajewa V. (1930): Infusorien des Salzigen Sees der Bucht Kruglaja in der Nähe von Sebastopol (Krim). – *Trudy sevastopol'. biol. Sta.* **2**: 31–46 (in Russian with German summary).
- Dalby A.B., Prescott D.M. (2004): The scrambled actin I gene in *Uroleptus pisces*. – *Chromosoma* **112**: 247–254.
- Deroux G. (1974): Quelques précisions sur *Strobilidium gyrans* Schewiakoff. – *Cah. Biol. mar.* **15**: 571–588.

- Detcheva R. (1977): Nouveaux ciliés de la mer Noire près de la côte bulgare. – *Acta Zool. Bulgarica* **8**: 3–6.
- Detcheva R.B. (1980): Composition et répartition des ciliés mésopsammiques des plages bulgares. – *Hydrobiology* **11**: 28–38.
- Detcheva R.B. (1981): Etude de la composition et de la répartition saisonnière des ciliés de la rivière Maritza. – *Hydrobiology* **14**: 16–30.
- Detcheva R.B. (1982): Recherches, en Bulgarie, sur les cilies de certaines plages de la Mer Noire et des rivières y aboutissant. – *Anns Stn limnol. Besse* **15** (years 1981/1982): 231–253.
- Detcheva R.B. (1983): Contribution sur la faune des ciliés mesopsammiques de quelques plages bulgares et celles des rivières qui se jettent dans la mer. – *Hydrobiology* **18**: 64–76.
- Detcheva R.B. (1983a): Caractéristiques écologiques des cilies de la riviere Maritza. – *Anns Stn limnol. Besse* **16** (years 1982/1983): 200–219.
- Detcheva R.B. (1992): *Catologi Faunae Bulgaricae 1 Protozoa, Ciliophora*. Academia Scientiarum Bulgaricae, Sofia. 134 pp.
- Diesing C.M. (1850): *Systema Helminthum* Vol. I. W. Braumüller, Vindobonae. I–XVI, 680 pp.
- Díaz S., Martín-González A., Borniquel S., Gutiérrez J.C. (2000): *Cyrtolophosis elongata* (Colpodea Ciliophora): some aspects of ciliary pattern, division, cortical and nuclear changes during encystment and resting cyst ultrastructure. – *Europ. J. Protistol.* **36**: 367–378.
- Dietz G. (1965): Über zwei neue Ciliaten aus einem ostafrikanischen Sodasee, *Uroleptus natronophilus* n. sp. und *Spathidium elmenteitanum* n. sp. – *Arch. Protistenk.* **108**: 25–28.
- Dini F., Nyberg D. (1993): Sex in ciliates. – *Adv. microb. Ecol.* **13**: 85–153.
- Dini F., Lucchesi P., Macchioni G. (1995): Protozoa. In: Minelli A., Ruffo S., La Posta S. (eds.): *Checklist delle specie della fauna italiana*, Calderini, Bologna. 92 pp.
- Dovgal I.V. (2002): Evolution, phylogeny and classification of Suctorea (Ciliophora). – *Protistology* **2**: 194–270.
- Dragesco J. (1953): Sur l'écologie des ciliés psammophiles littoraux de la région de Banyuls-sur-mer (Pyr.-Or.) (Note préliminaire) (I). – *Vie Milieu* **4**: 627–632.
- Dragesco J. (1954): Diagnoses préliminaires de quelques ciliés nouveaux des sables de Banyuls-sur-mer (I). – *Vie Milieu* **4**: 633–637.
- Dragesco J. (1954a): Diagnoses préliminaires de quelques ciliés nouveaux des sables. – *Bull. Soc. zool. Fr.* **79**: 62–70.
- Dragesco J. (1960): Ciliés mésopsammiques littoraux. Systématique, morphologie, écologie. – *Trav. Stn biol. Roscoff* **122**: 1–356.
- Dragesco J. (1962): L'orientation actuelle de la systématique des ciliés et la technique d'impregnation au protéinate d'argent. – *Bull. Micr. appl.* **11**: 49–58.
- Dragesco J. (1963): Compléments a la connaissance des ciliés mésopsammiques de Roscoff. – *Cah. Biol. mar.* **4**: 251–275.
- Dragesco J., Dragesco-Kernéis A. (1986): Ciliés libres de l'Afrique intertropicale. Introduction à la connaissance et à l'étude des ciliés. – *Faune tropicale (Éditions de l'ORSTOM)* **26**: 1–559.
- Dragesco J., Njine T. (1971): Compléments à la connaissance des ciliés libres du Cameroun. – *Anns Fac. Sci. Univ. féd. Cameroun* **7–8**: 97–140, Planches I, II.
- Dujardin F. (1841): *Histoire naturelle des zoophytes. Infusoires, comprenant la physiologie et la classification de ces animaux, et la manière de les étudier a l'aide du microscope*. Librairie Encyclopédique de Roret, Paris. I–XII, 684 pp, 14 pp Explication des Planches, Planches 1–22.
- Dzhurtubayev M.M. (1978): Species composition and vertical distribution of psammophilic infusoria in Odessa Bay. – *Hydrobiol. J.* **13**: 64–67.
- Ehrenberg C.G. (1830): Beiträge zur Kenntniß der Organisation der Infusorien und ihrer geographischen Verbreitung, besonders in Sibirien. – *Abh. preuss. Akad. Wiss., Phys.-math. Kl.* **year 1830**: 1–88, Tafeln I–VIII (Corliss 1979, p. 371, gives following comment to this reference: “Although this particular publication did not appear under this exact title, etc. until 1832, its identical content had, in effect, been available in distributable form since 1830, so I am taking the liberty of so dating the work for historical

- and nomenclatural purposes and usage". The title of the World List deviates somewhat from the original title which is "Physikalische Abhandlungen der Königlichen Akademie der Wissenschaften zu Berlin").
- Ehrenberg C.G. (1831): Über die Entwicklung und Lebensdauer der Infusionsthier; nebst ferneren Beiträgen zu einer Vergleichung ihrer organischen Systeme. – Abh. preuss. Akad. Wiss., Phys.- math. Kl. **year 1831**: 1–154, Tafeln I–IV (see note following entry on Ehrenberg 1830; similar situation, with "final" appearance again in 1832, but with initially availability in 1831 in this case).
- Ehrenberg C.G. (1833): Dritter Beitrag zur Erkenntniß großer Organisation in der Richtung des kleinsten Raumes. – Abh. preuss. Akad. Wiss., Phys.-math. Kl. **year 1833**: 145–336, Tafeln I–XI (See note following entry on Ehrenberg 1830; similar situation, except "final" appearance in 1835, in this case, but with initial availability in 1833).
- Ehrenberg C.G. (1838): Die Infusionsthierchen als vollkommene Organismen. Ein Blick in das tiefere organische Leben der Natur. L. Voss, Leipzig. 548 pp, Tafeln I–LXIV.
- Eigner P. (1995): Divisional morphogenesis in *Deviata abbrevescens* nov. gen., nov. spec., *Neogeneia hortualis* nov. gen., nov. spec., and *Kahliella simplex* (Horvath) Corliss and redefinition of the Kahliellidae (Ciliophora, Hypotrichida). – Europ. J. Protistol. **31**: 341–366.
- Eigner P. (1997): Evolution of morphogenetic processes in the Orthoamphisiellidae n. fam., Oxytrichidae, and Parakahliellidae n. fam., and their depiction using a computer method (Ciliophora, Hypotrichida). – J. Euk. Microbiol. **44**: 553–573.
- Eigner P. (1999): Comparison of divisional morphogenesis in four morphologically different clones of the genus *Gonostomum* and update of the natural hypotrich system (Ciliophora, Hypotrichida). – Europ. J. Protistol. **35**: 34–48.
- Eigner P. (2001): Divisional morphogenesis in *Uroleptus caudatus* (Stokes, 1886), and the relationship between the Urostylidae and the Parakahliellidae, Oxytrichidae, and Orthoamphisiellidae on the basis of morphogenetic processes (Ciliophora, Hypotrichida). – J. Euk. Microbiol. **48**: 70–79.
- Eigner P., Foissner W. (1991): *Orthoamphisiella stramenticola* nov. gen., nov. spec., a new hypotrichous ciliate (Ciliophora: Hypotrichida) occurring in walnut leaf litter. – Acta Protozool. **30**: 129–133.
- Eigner P., Foissner W. (1993): Divisional morphogenesis in *Orthoamphisiella stramenticola* Eigner & Foissner, 1991 and *O. grelli* nov. spec. (Ciliophora, Hypotrichida). – Arch. Protistenk. **143**: 337–345.
- Eigner P., Foissner W. (1994): Divisional morphogenesis in *Amphisiellides illuvialis* n. sp., *Paramphisiella caudata* (Hemberger) and *Hemiamphisiella terricola* Foissner, and redefinition of the Amphisiellidae (Ciliophora, Hypotrichida). – J. Euk. Microbiol. **41**: 243–261.
- Engelmann T.W. (1862): Zur Naturgeschichte der Infusionsthier. – Z. wiss. Zool. **11**: 347–393, Tafeln XXVIII–XXXI.
- Esteban G.F., Finlay B.J., Olmo J.L., Tyler P.A. (2000): Ciliated protozoa from a volcanic crater-lake in Victoria, Australia. – J. nat. Hist. (London) **34**: 159–189.
- Esteban G.F., Clarke K.J., Olmo J.L., Finlay B.J. (2006): Soil protozoa—An intensive study of population dynamics and community structure in an upland grassland. – Appl. Soil. Ecol. **33**: 137–151.
- Fanti L., Pimpinelli S. (1999): The peculiar organization of telomeres in *Drosophila melanogaster*. – Gene Therapy and Molecular Biology **4**: 1–10.
- Faria J.G.de, Cunha A.M.Da, Pinto C. (1922): Estudos sobre protozoarios do mar (Studien ueber Meeresprotozoen). – Mems Inst. Oswaldo Cruz **15**: 186–208, 101–115 (German text), Estampas 23–25.
- Fauré-Fremiet E. (1950): Écologie des ciliés psammophiles littoraux. – Bull. biol. Fr. Belg. **84**: 35–75.
- Fauré-Fremiet E. (1954): *Amphisiella lithophora*, n. sp., cilié hypotriche psammobie. – Bull. Soc. zool. Fr. **79**: 473–479.
- Fauré-Fremiet E. (1961): Remarques sur la morphologie comparée et la systématique des ciliata Hypotrichida. – C. r. hebd. Séanc. Acad. Sci., Paris **252**: 3515–3519.
- Fenchel T. (1968): The ecology of marine microbenthos II. The food of marine benthic ciliates. – Ophelia **5**: 73–121.
- Fenchel T. (1969): The ecology of marine microbenthos IV. Structure and function of the benthic ecosystem, its chemical and physical factors and the microfauna communities with special reference to the ciliated protozoa. – Ophelia **6**: 1–182.

- Fenchel T., Finlay B.J. (1995): Ecology and evolution in anoxic worlds. Oxford University Press, Oxford, New York, Tokyo. I–XII, 276 pp.
- Fenchel T., Jansson B-O. (1966): On the vertical distribution of the microfauna in the sediments of a brackish-water beach. – *Ophelia* **3**: 161–177.
- Fernández-Galiano D., Calvo P. (1992): Redescription of *Phacodinium metchnikoffi* (Ciliophora, Hypotrichida): general morphology and taxonomic position. – *J. Protozool.* **39**: 443–448.
- Fernandez-Leborans G. (1984): Description of *Amphisiella oscensis* sp. nov. (Protozoa: Ciliophora). – *J. nat. Hist. (London)* **18**: 25–30.
- Fernandez-Leborans G. (1985): The paroral formation of various hypotrichous ciliates. – *Arch. Protistenk.* **130**: 367–380.
- Fernandez-Leborans G. (2000): Protozoan species in three epibenthic areas of the Cantabrian Sea: relations with environmental actors. – *J. mar. biol. Ass. U. K.* **80**: 407–418.
- Fernandez-Leborans G. (2001): Relative importance of protozoan functional groups in three marine sublittoral areas. – *J. mar. biol. Ass. U. K.* **81**: 735–750.
- Fernandez-Leborans G., Antonio-García M.T. (1988): Effects of lead and cadmium in a community of protozoans. – *Acta Protozool.* **27**: 141–159.
- Fernandez-Leborans G., Novillo A. (1992): Morphology and taxonomy of two new species of marine ciliates (Ciliophora: Spirotrichea: Stichotrichida: Amphisiellidae). – *Proc. biol. Soc. Wash.* **105**: 165–179.
- Fernandez-Leborans G., Novillo A. (1993): Sublittoral protistan communities of the shores of the sea of Cantabria (Bay of Biscay). – *Int. Revue ges. Hydrobiol. Hydrogr.* **78**: 201–218.
- Fernandez-Leborans G., Novillo A. (1994): Effects of periodic addition of lead on a marine protistan community. – *Aquatic Sciences* **56**: 191–205.
- Fernandez-Leborans G., Novillo A. (2001): A note about two hypotrich ciliate species of the genus *Amphisiella*. – *Acta Protozool.* **40**: 225–227.
- Fernandez-Leborans G., Castro de Zaldumbide M., Perales C. (1990): Biomass and functional distribution according to the feeding modes of two freshwater protozoan communities. – *Int. Revue ges. Hydrobiol. Hydrogr.* **75**: 507–532.
- Fernandez-Leborans G., Valganon B., Castro de Zaldumbide M. (1999): Characterization of a marine sublittoral area facing the open sea, using epibenthic protists. – *Bull. mar. Sci.* **65**: 725–743.
- Finlay B.J., Rogerson A., Cowling A.J. (1988): A beginner's guide to the collection, isolation, cultivation and identification of freshwater protozoa. Culture Collection of Algae and Protozoa, Freshwater Biological Association, The Ferry House, Ambleside, Cumbria, United Kingdom. Titus Wilson and Son, Kendal, United Kingdom. 78 p.
- Finlay B.J., Esteban G.F., Olmo J.L., Tyler P.A. (1999): Global distribution of free-living microbial species. – *Ecography* **22**: 138–144.
- Finlay B.J., Esteban G.F., Clarke K.J., Olmo J.L. (2001): Biodiversity of terrestrial protozoa appears homogeneous across local and global spatial scales. – *Protist* **152**: 355–366.
- Fjeld P. (1955): On some marine psammobiotic ciliates from Drøbak (Norway) (with remarks on a method for quantitative studies on micropsammon). *Nytt magasin for zoologi* **3**: 5–59 (reference from Hartwig & Parker 1977).
- Fleury A., Iftode F., Deroux G., Fryd-Versavel G. (1985): Unité et diversité chez les hypotriches (Protozoaires ciliés): II – Éléments d'ultrastructure comparée chez divers représentants du sous-ordre des Euhypotrichina. – *Protistologica* **21**: 505–524.
- Fleury A., Iftode F., Deroux G., Fryd-Versavel G. (1986): Unité et diversité chez les hypotriches (Protozoaires ciliés): III. – Éléments d'ultrastructure comparée chez divers représentants dusous-ordre des Pseudohypotrichina et remarques générales. – *Protistologica* **22**: 65–87.
- Foissner W. (1976): *Wallackia schiffmanni* nov. gen., nov. spec. (Ciliophora, Hypotrichida) ein alpiner hypotricher Ciliat. – *Acta Protozool.* **15**: 387–392.
- Foissner W. (1979): Methylgrün-Pyronin: Seine Eignung zur supravitalen Übersichtsfärbung von Protozoen, besonders ihrer Protrichocysten. – *Mikroskopie* **35**: 108–115.

- Foissner W. (1981): Die Gemeinschaftsstruktur der Ciliatenzönose in alpinen Böden (Hohe Tauern, Österreich) und Grundlagen für eine Synökologie der terricolen Ciliaten (Protozoa, Ciliophora). – Veröff. Österr. MaB-Programms 4: 7–52.
- Foissner W. (1982): Ökologie und Taxonomie der Hypotrichida (Protozoa: Ciliophora) einiger österreichischer Böden. – Arch. Protistenk. **126**: 19–143.
- Foissner W. (1983a): Die Morphogenese von *Urosoma macrostyla* (Wrzesniowski, 1870) (Ciliophora: Oxytrichidae). – Arch. Protistenk. **127**: 413–428.
- Foissner W. (1983b): Morphologie und Morphogenese von *Psilotricha succisa* (O. F. Müller, 1786) nov. comb. (Ciliophora, Hypotrichida). – Protistologica **19**: 479–493.
- Foissner W. (1984): Infraciliatur, Silberliniensystem und Biometrie einiger neuer und wenig bekannter terrestrischer, limnischer und mariner Ciliaten (Protozoa: Ciliophora) aus den Klassen Kinetofragminophora, Colpodea und Polyhymenophora. – Stapfia **12**: 1–165.
- Foissner W. (1985): Protozoologische Untersuchungen an Almböden im Gasteiner Tal (Zentralalpen, Österreich). III. Struktur und Dynamik der Testaceen- und Ciliatentaxozönose. – Veröff. Österr. MaB-Programms **9**: 65–95.
- Foissner W. (1986): Beitrag zur Kenntnis der Bodenciliaten (Protozoa: Ciliophora) des Himalaja. – Zool. Jb. Syst. **113**: 45–53.
- Foissner W. (1986a): Wimpertiere (Protozoa: Ciliophora) in Flechten. – Katalog des Oberösterreichischen Landesmuseums Linz, Neue Folge **5**: 43–46.
- Foissner W. (1987): Soil protozoa: fundamental problems, ecological significance, adaptations in ciliates and testaceans, bioindicators, and guide to the literature. – Progr. Protistol. **2**: 69–212.
- Foissner W. (1987a): Neue und wenig bekannte hypotriche und colpodide Ciliaten (Protozoa: Ciliophora) aus Böden und Moosen. – Zool. Beitr. (N. F.) **31**: 187–282.
- Foissner W. (1987b): Faunistische und taxonomische Notizen über die Protozoen des Fuscher Tales (Salzburg, Österreich). – Jber. Haus Nat. Salzburg **10**: 56–68.
- Foissner W. (1988): Gemeinsame Arten in der terricolen Ciliatenfauna (Protozoa: Ciliophora) von Australien und Afrika. – Stapfia **17**: 85–133.
- Foissner W. (1991): Basic light and scanning electron microscopic methods for taxonomic studies of ciliated protozoa. – Europ. J. Protistol. **27**: 313–330.
- Foissner W. (1993): Colpodea (Ciliophora). – Protozoenfauna **4/1**: I–X + 1–798 pp.
- Foissner W. (1994): Morphology and morphogenesis of *Circinella arenicola* nov. gen., nov. spec., a cephalized hypotrich (Ciliophora, Hypotrichida) from sand dunes in Utah, USA. – Europ. J. Protistol. **30**: 156–170.
- Foissner W. (1994a): *Pentahymena corticicola* nov. gen., nov. spec., a new colpodid ciliate (Protozoa, Ciliophora) from bark of Acacia trees in Costa Rica. – Arch. Protistenk. **144**: 289–295.
- Foissner W. (1994b): Soil protozoa as bioindicators in ecosystems under human influence. In: Darbyshire J.F. (ed.): Soil protozoa, pp. 147–193. CAB International, Wallingford.
- Foissner W. (1994c): Progress in taxonomy of planktonic freshwater ciliates. – Marine Microbial Food Webs **8**: 9–35.
- Foissner W. (1995): Tropical protozoan diversity: 80 ciliate species (Protozoa, Ciliophora) in a soil sample from a tropical dry forest of Costa Rica, with descriptions of four new genera and seven new species. – Arch. Protistenk. **145**: 37–79.
- Foissner W. (1996): Ontogenesis in ciliated protozoa, with emphasis on stomatogenesis. In: Hausmann K., Bradbury P.C. (eds.): Ciliates: Cells as organisms, pp. 95–177. Gustav Fischer, Stuttgart Jena Lübeck Ulm.
- Foissner W. (1996a): Terrestrial ciliates (Protozoa, Ciliophora) from two islands (Gough, Marion) in the southern oceans, with description of two new species, *Arcuospathidium cooperi* and *Oxytricha ottowi*. – Biol. Fertil. Soils **23**: 282–291.
- Foissner W. (1996b): Faunistics, taxonomy and ecology of moss and soil ciliates (Protozoa, Ciliophora) from Antarctica, with description of new species, including *Pleuroplitoides smithi* gen. n., sp. n. – Acta Protozool. **35**: 95–123.
- Foissner W. (1997): Soil ciliates (Protozoa: Ciliophora) from evergreen rain forests of Australia, South America and Costa Rica: diversity and description of new species. – Biol. Fertil. Soils **25**: 317–339.

- Foissner W. (1997a): Soil ciliates (Protozoa: Ciliophora) from evergreen rain forests of Australia, South America and Costa Rica: diversity and description of new species. – *Biol. Fertil. Soils* **25**: 317–339.
- Foissner W. (1998): An updated compilation of world soil ciliates (Protozoa, Ciliophora), with ecological notes, new records, and descriptions of new species. – *Europ. J. Protistol.* **34**: 195–235.
- Foissner W. (1999): Notes on the soil ciliate biota (Protozoa, Ciliophora) from the Shimba Hills in Kenya (Africa): diversity and description of three new genera and ten new species. – *Biodiversity and Conservation* **8**: 319–389.
- Foissner W. (2000): Notes on ciliates (Protozoa, Ciliophora) from Espeletia trees and Espeletia soils of the Andean Páramo, with descriptions of *Sikorops espeletiae* nov. spec. and *Fragmocirrus espeletiae* nov. gen., nov. spec. – *Stud. Neotrop. Fauna & Environm* **35**: 52–79.
- Foissner W. (2000a): A compilation of soil and moss ciliates (Protozoa, Ciliophora) from Germany, with new records and descriptions of new and insufficiently known species. – *Europ. J. Protistol.* **36**: 253–283.
- Foissner W. (2002): Neotypification of protists, especially ciliates (Protozoa, Ciliophora). – *Bull. zool. Nom.* **59**: 165–169.
- Foissner W. (2004): Some new ciliates (Protozoa, Ciliophora) from an Austrian floodplain soil, including a giant, red “flagship”, *Cyrtohymena (Cyrtohymenides) aspoECKi* nov. subgen., nov. spec. – *Denisia* **13**: 369–382.
- Foissner W. (2005): The unusual, lepidosome-coated resting cyst of *Meseres corlissi* (Ciliophora: Oligotrichea): transmission electron microscopy and phylogeny. – *Acta Protozool.* **44**: 217–230.
- Foissner W. (2006): Biogeography and dispersal of micro-organisms: a review emphasizing protists. – *Acta Protozool.* **45**: 111–136.
- Foissner W., Adam H. (1983): Morphologie und Morphogenese des Bodenciliaten *Oxytricha granulifera* sp. n. (Ciliophora, Oxytrichidae). – *Zool. Scr.* **12**: 1–11.
- Foissner W., AL-Rasheid K.A.S. (2006): A unified organization of the stichotrichine oral apparatus, including a description of the buccal seal (Ciliophora: Spirotrichea). – *Acta Protozool.* **45**: 1–16.
- Foissner W., Blatterer H. (1990): New ciliates (Protozoa, Ciliophora) from Australia and Africa. – *J. Protozool. Suppl.* **37**: 9A, Abstract 53.
- Foissner W., Foissner I. (1988): Stamm: Ciliophora. – *Catalogus Faunae Austriae* **I** c: 1–147.
- Foissner W., Peer T. (1985): Protozoologische Untersuchungen an Almböden im Gasteiner Tal (Zentralalpen, Österreich). I. Charakteristik der Taxotope, Faunistik und Autökologie der Testacea und Ciliophora. – *Veröff. Österr. MaB-Programms* **9**: 27–50.
- Foissner W., Pichler M. (2006): The unusual, lepidosome-coated resting cyst of *Meseres corlissi* (Ciliophora: Oligotrichea): genesis of four complex types of wall precursors and assemblage of the cyst wall. – *Acta Protozool.* **45**: 339–366.
- Foissner W., Stoeck T. (2006): *Rigidothrix goiseri* nov. gen., nov. spec. (Rigidothrichidae nov. fam.), a new “flagship” ciliate from the Niger floodplain breaks the flexibility-dogma in the classification of stichotrichine spirotrichs (Ciliophora, Spirotrichea). – *Europ. J. Protistol.* **42**: 249–267.
- Foissner W., Xu K. (2007): Monograph of the Spathidiida (Ciliophora, Haptoria) Vol I: Protopathidiidae, Arcuospathidiidae, Apertospathulidae. – *Monographiae biol.* **81**: i–xi, 1–485.
- Foissner W., Franz H., Adam H. (1982): Terrestrische Protozoen als Bioindikatoren im Boden einer planierten Ski-Piste. – *Pedobiologia* **24**: 45–56.
- Foissner W., Peer T., Adam H. (1985): Pedologische und protozoologische Untersuchung einiger Böden des Tullnerfeldes (Niederösterreich). – *Mitt. öst. bodenk. Ges.* **30**: 77–117.
- Foissner W., Schlegel M., Prescott D.M. (1987): Morphology and morphogenesis of *Onychodromus quadricornutus* n. sp. (Ciliophora, Hypotrichida), an extraordinarily large ciliate with dorsal horns. – *J. Protozool.* **34**: 150–159.
- Foissner W., Franz H., Adam H. (1987a): Untersuchungen über das Bodenleben in ökologisch und konventionell bewirtschafteten Acker- und Grünlandböden in Raum Salzburg. – *Verh. Ges. Ökologie* **15** (year 1985): 333–339.
- Foissner W., Buchgraber K., Berger H. (1990): Bodenfauna, Vegetation und Ertrag bei ökologisch und konventionell bewirtschaftetem Grünland: Eine Feldstudie mit randomisierten Blöcken. – *Mitt. öst. bodenk. Ges.* **41**: 5–33.

- Foissner W., Oleksiv I., Müller H. (1990a): Morphologie und Infraciliatur einiger Ciliaten (Protozoa: Ciliophora) aus stagnierenden Gewässern. – Arch. Protistenk. **138**: 191–206.
- Foissner W., Blatterer H., Berger H., Kohmann F. (1991): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems – Band I: Cyrtophorida, Oligotrichida, Hypotrichida, Colpodea. – Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft **1/91**: 1–478.
- Foissner W., Berger H., Kohmann F. (1992): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems - Band II: Peritrichia, Heterotrichida, Odontostomatida. – Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft **5/92**: 1–502.
- Foissner W., Berger H., Blatterer H., Kohmann F. (1995): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems - Band IV: Gymnostomatea, *Loxodes*, Suctorina. – Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft **1/95**: 1–540.
- Foissner W., Berger H., Schaumburg J. (1999): Identification and ecology of limnetic plankton ciliates. – Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft **3/99**: 1–793.
- Foissner W., Agatha S., Berger H. (2002): Soil ciliates (Protozoa, Ciliophora) from Namibia (Southwest Africa), with emphasis on two contrasting environments, the Etosha region and the Namib desert. Part I: Text and line drawings. Part II: Photographs. – *Denisia* **5**: 1–1459.
- Foissner W., Moon-van der Staay S.Y., van der Staay G.W., Hackstein J.H.P., Krautgartner W.-D., Berger H. (2004): Reconciling classical and molecular phylogenies in the stichotrichines (Ciliophora, Spirotrichea), including new sequences from some rare species. – *Europ. J. Protistol.* **40**: 265–281.
- Foissner W., Berger H., Xu K., Zechmeister-Boltenstern S. (2005): A huge, undescribed soil ciliate (Protozoa: Ciliophora) diversity in natural forest stands of Central Europe. – *Biodiversity and Conservation* **14**: 617–701.
- Foissner W., Müller H., Weisse T. (2005a): The unusual, lepidosome-coated resting cyst of *Meseres corlissi* (Ciliophora: Oligotrichia): light and scanning electron microscopy, cytochemistry. – *Acta Protozool.* **44**: 201–215.
- Foissner W., Pichler M., Al-Rasheid K., Weisse T. (2006): The unusual, lepidosome-coated resting cyst of *Meseres corlissi* (Ciliophora: Oligotrichia): encystment and genesis and release of the lepidosomes. – *Acta Protozool.* **45**: 323–338.
- Foissner W., Müller H., Agatha S. (2007): A comparative fine structural and phylogenetic analysis of resting cysts in oligotrich and hypotrich Spirotrichea (Ciliophora). – *Europ. J. Protistol.* **43**: 295–314.
- Fromentel E.de (1874): Études sur les microzoaires ou infusoires proprement dits comprenant de nouvelles recherches sur leur organisation, leur classification et la description des espèces nouvelles ou peu connues. G. Masson, Paris. pp. I–VIII, 1–88.
- Froud J. (1949): Observations on hypotrichous ciliates: the genera *Stichotricha* and *Chaetospira*. – *Q. Jl microsc. Soc.* **90**: 141–158.
- Funke W. (1986): Tiergesellschaften im Ökosystem “Fichtenforst” (Protozoa, Metazoa-Invertebrata) - Indikatoren von Veränderungen in Waldökosystemen. – Kernforschungszentrum Karlsruhe GmbH, KfK-PEF **9**: 1–150.
- Gaievskaja N. (1925): Sur deux nouveaux infusoires des mares salées – *Cladotricha koltzowii* nov. gen. nov. sp. et *Palmarium salinum* nov. gen. nov. sp. – *Russk. Arkh. Protist.* **4**: 255–288, Planches XV, XVI.
- Galtsoff P.S., Lutz F.E., Welch P.S., Needham J.G. (1959): Culture methods for invertebrate animals. Dover Publications, New York (on the cover the authors are arranged as follows: Lutz, Welch, Galtsoff, Needham. On the spine only Needham is cited). I–XXXII, 590 pp.
- Ganner B., Unterweger A., Jäger P. (2002): Die Biologie der Salzburger Kläranlagen im Zeitraum von 1991 bis 2000. Zur Evaluation der Beurteilungskriterien bei der mikroskopischen Belebtschlammuntersuchung kommunaler Kläranlagen mit Stickstoffelimination. – *Gewässerschutz* **6**: 115–170.
- Gelei J. (1954): Über die Lebensgemeinschaft einiger temporärer Tümpel auf einer Bergwiese im Börzsönygebirge (Oberungarn) III. Ciliaten. *Acta biol. hung.* **5**: 259–343.
- Gellért J. (1956a): Die Ciliaten des sich unter der Flechte *Parmelia saxatilis* Mass. gebildeten Humus. – *Acta biol. hung.* **6**: 77–111.

- Gellért J. (1956b): Ciliaten des sich unter dem Moosrasen auf Felsen gebildeten Humus. – Acta biol. hung. **6**: 337–359.
- Gellért J. (1957): Néhány hazai lomblevelű és tűlevelű erdő talajának ciliáta-faunája Ciliatenfauna im Humus einiger ungarischen Laub- und Nadelholzwälder. – Anns Inst. biol. Tihany **24**: 11–34 (in Hungarian with German summary).
- Giambelluca M.A., Gabrielli S., Erra F., Rosati G. (1995): Morphogenetic study during cell division in *Euplotidium itoi* (Ciliata, Hypotrichida). – Europ. J. Protistol. **31**: 286–291.
- Gong J., Song W., Li L., Shao C., Chen Z. (2006): A new investigation of the marine ciliate, *Trachelostyla pediculiformis* (Cohn, 1866) Borror, 1972 (Ciliophora, Hypotrichida), with establishment of a new genus, *Spirotrachelostyla* nov. gen. – Europ. J. Protistol. **42**: 63–73.
- Gong J., Kim S.-J., Kim S.-Y., Min G.-S., Roberts D. McL., Warren A., Choi J.-K. (2007): Taxonomic re-descriptions of two ciliates, *Protoastrostyla pulchra* n. g., n. comb. and *Hemigastrostyla enigmatica* (Ciliophora: Spirotrichea, Stichotrichia), with phylogenetic analyses based on 18S and 28S rRNA gene sequences. – J. Euk. Microbiol. **54**: 468–478.
- Gourret P., Roeser P. (1888): Contribution à l'étude des protozoaires de la Corse. – Archs Biol. **8**: 139–204, Planches XIII–XV.
- Grandori R., Grandori L. (1934): Studi sui protozoi del terreno. – Boll. Lab. Zool. agr. Bachic. R. Ist. sup. agr. Milano **5**: 1–341, Tavole I–XIV.
- Griffin L.E. (1910): *Euplotes worcesteri* sp. nov.: II. Division. – Philipp. J. Sci., D. Ethnology, Anthropology, and General Biology **5**: 315–336, Plate IV–VIII.
- Grim J.N., Manganaro C.A. (1985): Form of the extrusomes and secreted material of the ciliated protozoan *Pseudourostyla cristata*, with some phylogenetic interpretations: a light, scanning electron, and transmission electron microscopic study. – Trans. am. microsc. Soc. **104**: 350–359.
- Grispini M. (1938): I ciliati di una pozza di «macchia» della campagna romana. – Boll. Zool. **9**: 147–158.
- Grolière C.-A. (1975): Descriptions de quelques ciliés hypotriches des tourbières a sphaignes et des étendues d'eau acides. – Protistologica **11**: 481–498.
- Gruber A. (1884): Die Protozoen des Hafens von Genua. – Nova Acta Acad. Caesar. Leop. Carol. **46**: 473–539, Tafeln VII–XI.
- Gruber A. (1888): Weitere Beobachtungen an vielkernigen Infusorien. – Ber. naturf. Ges. Freiburg i. B. **3**: 57–70, Tafeln VI, VII.
- Gupta R., Kamra K., Sapra G.R. (2006): Morphology and cell division of the oxytrichids *Architricha indica* nov. gen., nov. sp., and *Histiculus histrio* (Müller, 1773), Corliss, 1960 (Ciliophora, Hypotrichida). – Europ. J. Protistol. **42**: 29–48.
- Gutiérrez J.C., Díaz S., Ortega R., Martin-González A. (2003): Ciliate resting cyst walls: a comparative review. – Recent Res. Devel. Microbiology **7**: 361–379.
- H**amburger C., Buddenbrock W.von (1929): Nordische Ciliata mit Ausschluss der Tintinoidea. – Nord. Plankt. **7**: 1–152.
- Hartwig E. (1973): Die Ciliaten des Gezeiten-Sandstrandes der Nordseeinsel Sylt I. Systematik. – Abh. math.-naturw. Kl. Akad. Wiss. Mainz, Reihe Mikrofauna des Meeresbodens **18**: 385–453.
- Hartwig E. (1973a): Die Ciliaten des Gezeiten-Sandstrandes der Nordseeinsel Sylt II. Ökologie. – Abh. math.-naturw. Kl. Akad. Wiss. Mainz, Reihe Mikrofauna des Meeresbodens **21**: 1–171.
- Hartwig E. (1973b): Die Nahrung der Wimpertiere des Sandlückensystems. – Mikrokosmos **62**: 329–336.
- Hartwig E. (1974): Verzeichnis der im Bereich der deutschen Meeresküste angetroffenen interstitiellen Ciliaten. – Mitt. hamb. zool. Mus. Inst. **71**: 7–21.
- Hartwig E. (1980): The marine interstitial ciliates of Bermuda with notes on their geographical distribution and habitat (1). – Cah. Biol. mar. **21**: 409–441.
- Hartwig E. (1984): Experimentelle Untersuchungen zur Wirkung von Rohöl und Rohöl/Tensid-Gemischen im Ökosystem Wattenmeer. IX. Benthische Ciliaten (Protozoa). – Senckenberg. marit. **16**: 121–151.
- Hartwig E., Parker J.G. (1977): On the systematics and ecology of interstitial ciliates of sandy beaches in North Yorkshire. – J. mar. biol. Ass. U. K. **57**: 735–760.

- Hausmann K., Bradbury P.C. (eds.) (1996): Ciliates. Cells as organisms. Gustav Fischer, Stuttgart, Jena, Lübeck, Ulm. i–x, 485 pp.
- Hausmann K., Hülsmann N. (1996): Protozoology. Georg Thieme Verlag, Stuttgart New York (with contributions by H. Macheimer, M. Mulisch, and G. Steinbrück). I–VIII, 338 pp.
- Hausmann K., Hülsmann N. (1996a): "Einzellige Eukaryota", Einzeller. In: Westheide W., Rieger R. (eds.): Spezielle Zoologie. Erster Teil: Einzeller und Wirbellose Tiere, pp. 1–72. Gustav Fischer Verlag, Stuttgart, Jena, New York.
- Hausmann K., Hülsmann N., Radek R. (2003): Protozoology. E. Schweizerbart'sche Verlagsbuchhandlung, Berlin, Stuttgart. I–X, 379 pp.
- Hemberger H. (1982): Revision der Ordnung Hypotrichida Stein (Ciliophora, Protozoa) an Hand von Protargolpräparaten und Morphogenesedarstellungen. Dissertation Universität Bonn. I–IV, 296 pp.
- Hemberger H. (1985): Neue Gattungen und Arten hypotricher Ciliaten. – Arch. Protistenk. **130**: 397–417.
- Hentschel E.J., Wagner G.H. (1996): Zoologisches Wörterbuch. Tiernamen, allgemeinbiologische, anatomische, physiologische Termini und Kurzbiographien. Fischer Verlag, Jena. 677 pp.
- Hewitt E.A., Müller K.M., Cannone J., Hogan D.J., Gutell R., Prescott D.M. (2003): Phylogenetic relationships among 28 spirotrichous ciliates documented by rDNA. – Molecular Phylogenetics and Evolution **29**: 258–267.
- Hill B.F. (1990): *Uronychia transfuga* (O. F. Müller, 1786) Stein, 1859 (Ciliophora, Hypotrichida, Uronychiidae): cortical structure and morphogenesis during division. – J. Protozool. **37**: 99–107.
- Hill B.F., Borror A.C. (1992): Redefinition of the genera *Diophrys* and *Paradiophrys* and establishment of the genus *Diophryopsis* n. g. (Ciliophora, Hypotrichida): implication for the species problem. – J. Protozool. **39**: 144–153.
- Hogan D.J., Hewitt E.A., Orr K.E., Prescott D.M., Müller K.M. (2001): Evolution of IESs and scrambling in the actin I gene in hypotrichous ciliates. – Proc. natn. Acad. Sci. U.S.A. **98**: 15101–15106.
- Horváth J.v. (1932): Ein neues hypotriches Infusor, *Kahlia acrobates* nov. gen., nov. sp. – Arch. Protistenk. **77**: 424–433.
- Horváth J.v. (1933): Beiträge zur hypotrichen Fauna der Umgebung von Szeged. I. – Arch. Protistenk. **80**: 281–302.
- Horváth J. (1950): Contributions to studies on soil protozoa of the ciliata group, with special regard to their adaptation to soil conditions. – Anns Inst. Biol. pervest. hung. **19** (years 1949–1950): 151–162.
- Hu X., Song W. (2000): Morphology and morphogenesis of a marine ciliate, *Gastrostyla pulchra* (Perejaslawzewa, 1885) Kahl, 1932 (Ciliophora, Hypotrichida). – Europ. J. Protistol. **36**: 201–210.
- Hu X., Song W. (2002): Studies on the ectocommensal ciliate, *Trachelostyla tani* nov. spec. (Protozoa: Ciliophora: Hypotrichida) from the mantle cavity of the scallop *Chlamys farrei*. – Hydrobiologia **481**: 173–179.
- Hu X., Song W. (2003): Redescription of two known species, *Gastrocirrhus monilifer* (Ozaki et Yagiu, 1942) and *Gastrocirrhus stentoreus* Bullington, 1940, with reconsideration of the genera *Gastrocirrhus* and *Euplotidium*. – Acta Protozool. **42**: 345–355.
- Hu X., Gong J. Song W. (2003): Pathogenic ciliates in scallop-farming waters. In: Song W., Zhao Y., Xu K., Hu X., Gong J. (eds.): Pathogenic protozoa in mariculture, pp. 145–178. Science Press, Beijing (in Chinese).
- Hu X., Warren A., Sudzuki T. (2004): Morphology and morphogenesis of two marine ciliates, *Pseudokeronopsis pararubra* sp. n. and *Amphisiella annulata* from China and Japan (Protozoa: Ciliophora). – Acta Protozool. **43**: 351–368.
- ICZN (The international Commission on Zoological Nomenclature) (1964): International code of zoological nomenclature. Second edition adopted by the XV International Congress of Zoology. International Trust for Zoological Nomenclature, London I–XX, 176 pp.
- ICZN (The international Commission on Zoological Nomenclature) (1985): International code of zoological nomenclature. Third edition adopted by the XX General Assembly of the International Union of Biological Sciences. University of California Press, Berkeley Los Angeles. I–XX, 338 pp.

- ICZN (The International Commission on Zoological Nomenclature) (1999): International Code of zoological nomenclature. International Trust for Zoological Nomenclature, London. 306 pp.
- Ilowaisky S.A. (1921): Zwei neue Arten und Gattungen von Infusorien aus dem Wolgabassin. – *Rab. volzh. biol. Sta.* **6**: 97–105, Tafeln VI, VII (in Russian with detailed German summary).
- Jaeckel S.G.A. (1962): Die Tierwelt der Schlei. Übersicht einer Brackwasserfauna. – *Schr. naturw. Ver. Schlesw.-Holst.* **33**: 11–32.
- Jankowski A.W. (1975): A conspectus of the new system of subphylum Ciliophora Doflein, 1901. – *Akad. Nauk. SSSR, Zool. Inst. Leningrad year 1975*: 26–27.
- Jankowski A.W. (1978): The revision of the system of Polyhymenophora class (Spirotrichia). – *Tezisy Dokl. zool. Inst. Akad. Nauk SSSR year 1978*: 39–40 (in Russian; translation of title and source after *Zool. Rec.*, 116, Year 1979, p. 144; according to the Zoological Record, the spelling of the author is Yankovskij A.V.).
- Jankowski A.W. (1979): Revision of the order Hypotrichida Stein, 1859. Generic catalogue, phylogeny, taxonomy. – *Trudy zool. Inst., Leningr.* **86**: 48–85 (in Russian with English summary).
- Jankowski A.W. (1989): Replacement of unacceptable generic names in the phylum Ciliophora. – *Vestnik Zoologii 1989*: 86.
- Jersabek C.D., Schabetsberger R. (1996): Limnological aspects of an alpine karst lake with extreme changes in water level. – *Limnologica (Berlin)* **26**: 1–13.
- Jones E.E. (1974): The protozoa of Mobile Bay, Alabama. – *Univ. South Alabama Monogr.* **1**: 1–113.
- Kahl A. (1928a): Die Infusorien (Ciliata) der Oldesloer Salzwasserstellen. – *Arch. Hydrobiol.* **19**: 50–123.
- Kahl A. (1928b): Die Infusorien (Ciliata) der Oldesloer Salzwasserstellen. – *Arch. Hydrobiol.* **19**: 189–246.
- Kahl A. (1932): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 3. Spirotricha. – *Tierwelt Dtl.* **25**: 399–650.
- Kahl A. (1933): Ciliata libera et ectocommensalia. – *Tierwelt N.- u. Ostsee Lieferung 23 (Teil II. c<sub>3</sub>)*: 29–146.
- Kahl A. (1930): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 1. Allgemeiner Teil und Prostomata. – *Tierwelt Dtl.* **18**: 1–180.
- Kahl A. (1931): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 2. Holotricha außer den im 1. Teil behandelten Prostomata. – *Tierwelt Dtl.* **21**: 181–398.
- Kahl A. (1932): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 3. Spirotricha. – *Tierwelt Dtl.* **25**: 399–650.
- Kahl A. (1935): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 4. Peritricha und Chonotricha. – *Tierwelt Dtl.* **30**: 651–886, I–V.
- Kamra K., Sapra G.R. (1990): Partial retention of parental ciliature during morphogenesis of the ciliate *Coniculostomum monilata* (Dragesco & Njiné, 1971) Njiné, 1978 (Oxytrichidae, Hypotrichida). – *Europ. J. Protistol.* **25**: 264–278.
- Kattar M.R. (1970): Estudo dos protozoários ciliados psamófilos do litoral brasileiro. – *Zoologia e biologia marinha N. S. N° 27*: 123–206.
- Katz L.A., McManus G.B., Snoeyenbos-West O.L.O., Griffin A., Pirog K., Costas B., Foissner W. (2005): Reframing the ‘Everything is everywhere’ debate: evidence for high gene flow and diversity in ciliate morphospecies. – *Aquat. microb. Ecol.* **41**: 55–65.
- Kent W.S. (1880–1882): A manual of the infusoria: including a description of all known flagellate, ciliate, and tentaculiferous protozoa, British and foreign, and an account of the organization and affinities of the sponges. Volumes I–III. David Bogue, London (Vol. I 1880: 1–432; Vol. II 1881: 433–720; Vol. II 1882: 721–913; Vol. III 1882: Plates I–LI). 913 pp.
- Kerney M.P., Cameron R.A.D., Jungbluth J.H. (1983): Die Landschnecken Nord- und Mitteleuropas. Ein Bestimmungsbuch für Biologen und Naturfreunde. P. Parey, Hamburg Berlin. 384 pp.
- Kiesselbach A. (1936): Zur Ciliatenfauna der nördlichen Adria. – *Thalassia* **2**: 1–53, 8 Tafeln.

- Kirby H. (1945): The structure of the common intestinal trichomonad of man. – *J. Parasitol.* **31**: 163–174.
- Klein B.M. (1936): Beziehungen zwischen Maschenweite und Bildungsvorgängen im Silberliniensystem der Ciliaten. – *Arch. Protistenk.* **88**: 1–22, Tafel 1.
- Klobutcher L.A., Prescott D.M. (1986): The special case of the hypotrichs. In: Gall J.G. (ed.): *The molecular biology of ciliated protozoa*, pp. 111–154. Academic Press, Orlando, San Diego, New York, Austin, Boston, London, Sydney, Tokyo, Toronto. I–VIII, 352 pp.
- Klobutcher L.A., Swanton M.T., Donini P., Prescott D.M. (1981): All gene-sized DNA molecules in four species of hypotrichs have the same terminal sequence and an unusual 3' terminus. – *Proc. natn. Acad. Sci. U.S.A.* **78**: 3015–3019.
- Knoop V., Müller K. (2006): *Gene und Stammbäume. Ein Handbuch zur molekularen Phylogenetik.* Elsevier Spektrum Akademischer Verlag, München. I–VII, 310 pp.
- Kool R. (1970): Morphogenesis of *Trachelostyla pediculiformis* (Cohn, 1866) (Ciliophora, Hypotrichida). – *J. Protozool. Suppl.* **17**: 6, Abstract 1.
- Kovaleva V.G. (1966): Infusoria of the mesopsammon in sand bays of the Black Sea. – *Zool. Zh.* **45**: 1600–1611 (in Russian with English summary).
- Kovaleva V.G., Golemansky V.G. (1979): Psammobiotic ciliates of the Bulgarian coast of the Black Sea. – *Acta Protozool.* **18**: 265–284 (in Russian with English summary).
- Kovaljeva V.G. (1967): New data on the infusorian fauna of the mesopsammon of the Barentz Sea. – *Acta Protozool.* **5**: 81–88 (in Russian with English summary).
- Kowalewskiego M. (1882): Przyczynek do historyi naturalnej oxytrichów. – *Pam. fizyogr.* **2**: 395–413, Tablica XXIX, XXX (in Polish with French explanation of plates).
- Krainer K.H. (1999): Vorläufiges Verzeichnis der Wimpertierchen Kärntens (Ciliata). – *Naturschutz in Kärnten* **15**: 665–671.
- Küsters E. (1974): Ökologische und systematische Untersuchungen der Aufwuchsciliaten im Königshafen bei List/Sylt. – *Arch. Hydrobiol. Suppl.* **45**: 121–211.
- Lackey J.B. (1961): Bottom sampling and environmental niches. – *Limnol. Oceanogr.* **6**: 271–279.
- Lackey J.B., Lackey E.W. (1963): Microscopic algae and protozoa in the waters near Plymouth in August 1962. – *J. mar. biol. Ass. U. K.* **43**: 797–805.
- Lamouroux J.-V.-F., Bory de Saint-Vincent J.B., Deslongchamps E. (1824): *Encyclopédie méthodique. Histoire naturelle des zoophytes, ou animaux rayonnés, faisant suite a l'histoire naturelle des vers de bruguière.* Vol. II, Agasse, Paris (according to the “Catalogue général des livres imprimés de la bibliothèque nationale, Paris” 1824 is the correct year). I–VIII, 819 pp.
- Laval-Peuto M. (1975): Cortex, périlemme et réticulum vésiculeux de *Cyrtarocyelis brandti* (Cilié tintinnide). Les ciliés a périlemme. – *Protistologica* **11**: 83–98.
- Laval-Peuto M. (1994): Classe des Oligotrichea Bütschli, 1887. *Ordre des Tintinnida* Kofoid et a Campbell, 1929. – *Traite de Zoologie* **2**: 181–219.
- Lecointre G., Le Guyader H. (2001): *Biosystematik.* Springer, Berlin, Heidelberg, New York. I–XII, 696 pp.
- Lee J.J., Small E.B., Lynn D.H., Bovee E.C. (1985): Some techniques for collecting, cultivating and observing protozoa. In: Lee J.J., Hutner S.H., Bovee E.C. (eds.). *An illustrated guide to the protozoa*, pp. 1–7. Society of Protozoologists, Lawrence.
- Lehle E. (1989): Beiträge zur Fauna der Ulmer Region II. Ciliaten (Protozoa:Ciliophora) Bioindikatoren in Waldböden. – *Mitt. Ver. Naturwiss. Math. Ulm (Donau)* **35**: 131–156.
- Lehle E. (1993): Bodenciliaten (Protozoa: Ciliophora) als sensitive Indikatoren tiefgreifender Veränderungen in Waldökosystemen. – *Kernforschungszentrum Karlsruhe GmbH, KfK-PEF* **113**: 1–91.
- Lehle E. (1994): Die Auswirkungen von Düngung und Kalkung auf die Bodenciliaten (Protozoa: Ciliophora) eines Fichtenbestandes im Schwarzwald (Süddeutschland). – *Arch. Protistenk.* **144**: 113–125.
- Lehle E., Jans W., Funke W. (1992): Reaktionen von Bodenciliaten auf Düngemaßnahmen in Fichtenforsten des Schwarzwaldes. – *Kernforschungszentrum Karlsruhe GmbH, KfK-PEF* **94**: 273–287.

- Lei Y., Choi J.K., Xu K. (2002): Morphology and infraciliature of a new marine ciliate *Paraeuplotidium smalli* n. sp. with description of a new genus, *Paraeuplotidium* n. g. (Ciliophora, Euplotida). – J. Euk. Microbiol. **49**: 402–406.
- Lepsi J. (1928): Un nouveau protozoaire marin: *Gastrocirrhus intermedius*. – Anns Protist. **1**: 195–197.
- Lepsi I. (1962): Über einige insbesondere psammobionte Ciliaten vom rumänischen Schwarzmeer-Ufer. – Zool. Anz. **168**: 460–465.
- Li J., Lin X., Shao C., Gong J., Hu X., Song W. (2007): Morphological redescription and neotypification of the marine ciliate, *Amphisiella marioni* Gourret & Roeser, 1888 (Ciliophora: Hypotrichida), a poorly known form misidentified for a long time. – J. Euk. Microbiol. **54**: 364–370.
- Li L., Song W. (2006): Phylogenetic position of the marine ciliate, *Certesias quadrinucleata* (Ciliophora; Hypotrichia; Hypotrichida) inferred from the complete small subunit ribosomal RNA gene sequence. – Europ. J. Protistol. **42**: 55–61.
- Li L., Song W., Hu X. (2007): Two marine hypotrichs from north China, with description of *Spiroamphisiella hembergeri* gen. nov., spec. nov. (Ciliophora, Hypotricha). – Acta Protozool. **46**: 107–120.
- Lin X., Song W. (2004): Redescription of the marine ciliate, *Certesias quadrinucleata* (Protozoa: Ciliophora) from Qingdao, China. – J. mar. biol. Ass. U. K. **84**: 1131–1136.
- Lin X., Song W., Warren A. (2004): Redescription of the rare marine ciliate, *Prodiscocephalus borrori* (Wicklow, 1982) from shrimp-culturing waters near Qingdao, China, with redefinitions of the genera *Discocephalus*, *Prodiscocephalus* and *Marginotricha* (Ciliophora, Hypotrichida, Discocephalidae). – Europ. J. Protistol. **40**: 137–146.
- Lincoln R.J., Boxshall G.A., Clark P.F. (1985): A dictionary of ecology, evolution and systematics. Cambridge University Press, Cambridge, London, New York, New Rochelle, Melbourne, Sydney. 298 pp.
- Lüftenecker G., Foissner W., Adam H. (1986): Der Einfluss organischer und mineralischer Dünger auf die Bodenfauna einer planierten, begrüneten Schipiste oberhalb der Waldgrenze. – Z. Vegetationstechnik **9**: 149–153.
- Lüftenecker G., Foissner W., Berger H., Adam H. (1987): Ciliatostasis and its disruption by human influences. – Society of Protozoologists **1986 Abstracts**: Abstract 139.
- Lüftenecker G., Petz W., Foissner W., Adam H. (1988): The efficiency of a direct counting method in estimating the numbers of microscopic soil organisms. – Pedobiologia **31**: 95–101.
- Lynn D.H. (1991): The implications of recent descriptions of kinetid structure to the systematics of the ciliated protists. – Protoplasma **164**: 123–142.
- Lynn D.H. (2003): Morphology or molecules: how do we identify the major lineages of ciliates (Phylum Ciliophora)? – Europ. J. Protistol. **39**: 356–364.
- Lynn D.H., Corliss J.O. (1991): Ciliophora. In: Harrison F.W. (ed.). Microscopic anatomy of invertebrates Volume 1: Protozoa, pp. 333–467. Wiley-Liss, New York, Chichester.
- Lynn D.H., Small E.B. (2002): Phylum Ciliophora. In: Lee J.J., Leedale G.F., Bradbury P. (eds.): An illustrated guide to the Protozoa. Second edition. Organisms traditionally referred to as Protozoa, or newly discovered groups. Volume I, pp. 371–656. Society of Protozoologists, Lawrence.
- Madoni P. (2006): Benthic ciliates in Adriatic Sea lagoons. – Europ. J. Protistol. **42**: 165–173.
- Madrazo-Garibay M., López-Ochoterena E. (1982): Segunda lista taxonómica comentada de protozoarios de vida libre de México. – Revta lat.-am. Microbiol. **24**: 281–295.
- Madrazo-Garibay M., López-Ochoterena E. (1985): Protozoarios ciliados de México. XXVI. Análisis morfológico y taxonómico de treinta y cinco especies de la Laguna de Términos, Campeche. – Anales Instituto de Ciencias del Mar y Limnología Universidad Nacional Autonomía de México **12**: 199–212, Láminas 1, 2.
- Maeda M., Carey P.G. (1984): A revision of the genera *Trachelostyla* and *Gonostomum* (Ciliophora, Hypotrichida), including redescriptions of *T. pediculiformis* (Cohn, 1866) Kahl, 1932 and *T. caudata* Kahl, 1932. – Bull. Br. Mus. nat. Hist. (Zool.) **47**: 1–17.
- Mansfeld K. (1923): 16 neue oder wenig bekannte marine Infusorien. – Arch. Protistenk. **46**: 97–140.

- Martin J., Fedriani C., Nieto J. (1981): Étude comparée des processus morphogénétiques d'*Uroleptus* sp. (Kahl, 1932) et de *Holosticha* (*Paruroleptus*) *musculus* (Kahl, 1932) (Ciliés hypotriches). – *Protistologica* **17**: 215–224.
- Martin-Gonzalez A., Benit L., Palacios G., Gutierrez J.C. (1992): Ultrastructural analysis in resting cysts and encystment in *Colpoda inflata* I Normal and abnormal resting cysts. – *Cytobios* **72**: 7–18.
- Martinez E.A. (1980): Sensitivity of marine ciliates (Protozoa, Ciliophora) to high thermal stress. – *Estuar. Coastal Mar. Sci.* **10**: 369–381.
- Martínez-Murillo M.E., Aladro-Lubel M.A. (1994): Ciliados asociados al pasto marino *Halodule beaudettei* en la laguna de Tamiahua, Veracruz, México. – *An. Inst. Biol. Univ. Méx.* **65**: 11–18.
- Matis D., Tirjaková E., Stloukal E. (1996): Nálevníky (Ciliophora) v databanke fauny Slovenska (Ciliophora in the database of Slovak fauna). – *Folia Faunistica Slovaca* **1**: 3–38.
- Matsusaka T., Noguchi O., Yonezawa F. (1989): Cortical morphogenesis during encystment in a ciliate, *Euplotes encysticus* Yonezawa, 1985. – *Europ. J. Protistol.* **24**: 133–137.
- Matthes D. (1988): Suctoria (Sauginfusorien). – *Protozoenfauna* **7/1**: I–XIII, 1–226, 307–309.
- Maupas E. (1883): Contribution a l'étude morphologique et anatomique des infusoires ciliés. – *Archs Zool. exp. gén.*, 2<sup>e</sup> Série **1**: 427–664, Planches XIX–XXIV.
- Mayén Estrada R., Aladro Lubel M.A. (1987): Treinta especies de protozoarios ciliados bentonicos de Laguna de la Mancha, Veracruz. – *Universidad y Ciencia* **4**: 69–80.
- Mayer M. (1981): Kultur und Präparation der Protozoen. Franckh'sche Verlagshandlung, Stuttgart. 83 pp.
- Mazei Y.A., Burkovskii I.V., Stolyarov A.P. (2002): Salinity as a factor of forming ciliate community (Colonization experiments). – *Zool. Zh.* **81**: 387–393 (in Russian with English summary).
- Mazei Y.A., Burkovsky I.V. (2003): Vertical structure of the interstitial ciliate community in the Chernaya river estuary (the White Sea). – *Protistology* **3**: 107–120.
- Mazei Y.A., Burkovsky I.V. (2005): Species composition of benthic ciliate community in the Chernaya River estuary (Kandalaksha Bay, White Sea) with a total checklist of the White Sea benthic ciliate fauna. – *Protistology* **4**: 107–120.
- Mazei Y.A., Burkovsky I.V. (2006): Patterns of psammophilous ciliate community structure along the salinity gradient in the Chernaya River estuary (the Kandalaksha Gulf, the White Sea). – *Protistology* **4**: 251–268.
- Mereschkowsky C.von (1877): Studien über Protozoen des nördlichen Russland. – *Trudy imp. S-peterb. Obshch. Estest.* **8**: 203–385, Tafeln I–III (in Russian; translation of authors name and title from Mereschkowsky 1879).
- Miao M., Song W., Chen Z., AL-Rasheid K.A.S., Shao C., J. Jiang, Guo W. (2007): A unique euplotid ciliate, *Gastrocirrhus* (Protozoa, Ciliophora): assessment of its phylogenetic position inferred from the small subunit rRNA gene sequence. – *J. Euk. Microbiol.* **54**: 371–378.
- Milne W. (1886): On a new tentaculiferous protozoon and other infusoria, with notes on reproduction and the function of the contractile vesicle. – *Proc. Manchr Fld Nat. Archaeol. Soc.* **18**: 48–56, Plate I (Usually cited as “Proceedings. Philosophical Society of Glasgow”. However, according to the World List the other name is correct).
- Miyake A. (1996): Fertilization and sexuality in ciliates. In: Hausmann K., Bradbury P.C. (eds.): *Ciliates. Cells as organisms*, pp. 243–290. Gustav Fischer Verlag, Stuttgart, Jena, Lübeck, Ulm.
- Modeo L., Petroni G., Rosati G., Montagnes D.J.S. (2003): A multidisciplinary approach to describe protists: redescrptions of *Novistrombidium testaceum* Anigstein 1914 and *Strombidium inclinatum* Montagnes, Taylor, and Lynn 1990 (Ciliophora, Oligotrichia). – *J. Euk. Microbiol.* **50**: 175–189.
- Moskowitz N. (1950): The use of protein silver for staining protozoa. – *Stain Technol.* **25**: 17–20.
- Müller H. (2007): Live observation of excystment in the spirotricha ciliate *Meseres corlissi*. – *Europ. J. Protistol.* **43**: 95–100.
- Müller O.F. (1773): Vermium Terrestrium et Fluviatilium, seu Animalium Infusorium, Helminthicorum et Testaceorum, non Marinorum, Succincta Historia. Heineck & Faber, Havniae & Lipsiae. 135 pp.
- Münch H.-D. (1956): Spirotriche und peritriche Ciliaten, Rotatorien und Archianneliden aus dem Küstengrundwasser. (Anhang: Oligochaeten und Copepoden). – *Wiss. Z. Ernst Moritz Arndt-Univ. Greifswald, Mathematisch-naturwissenschaftliche Reihe* **5**: 433–436.

- Myskova O.M. (1976): Supplement to the fauna of psammophilous infusoria from Peter the Great Bay (Sea of Japan). – Sb. Rab. Inst. biol. Morya **5**: 83–86 (in Russian with English summary).
- Niebuhr J.W.H. (1989): Ökologische Studien an Bodenprotozoenpopulationen (Protozoa: Ciliophora) auf dem Gelände eines ehemaligen Textilindustriebetriebes in Nordhorn in Bezug auf ihre Verwendung als Bio-Indikatororganismen für anthropogen belastete Böden. – unter Berücksichtigung einiger Verfahren aus der Spurenanalytik. Diplomarbeit Universität Oldenburg. 181 pp.
- Nixon K.C., Wheeler Q.-D. (1990): An amplification on the phylogenetic species concept. – *Cladistics* **6**: 211–223.
- Njine T. (1978): Contribution a l'étude des ciliés libres du Cameroun. *Ecologie - cytologie*. Thèse Université Clermont- Ferrand II. 201 pp.
- Njiné T. (1979): Compléments a l'étude des ciliés libres du Cameroun. – *Protistologica* **15**: 343–354.
- Nobili R. (1957): Contributo all'ecologia dei ciliati psammofili del Golfo di Napoli. – *Boll. Zool.* **24**: 211–225.
- Nobili R., Rosati Raffaelli G. (1971): The fine structure of the oral apparatus of some ciliate hypotrichs. – *Monitore zool. ital., (N. S.)* **5**: 65–80.
- O**lins D.E., Olins A.L. (1994): The replication band of ciliated protozoa. – *Int. Rev. Cytol.* **153**: 137–170.
- Olmo Rísquez J.L. (1998): Diversidad local y global de los protozoos ciliados de hábitats de agua dulce. Tesis Universidad Complutense de Madrid, Madrid. 108 pp.
- P**ackroff G., Zwick P. (1996): The ciliate fauna of an unpolluted foothill stream, the Breitenbach, 1: Qualitative aspects. – *Limnologica (Berlin)* **26**: 255–262.
- Paiva T. da S., Silva-Neto I.D. da (2007): Morphology and morphogenesis of *Strongylidium pseudocrasum* Wang and Nie, 1935, with redefinition of *Strongylidium* Sterki, 1878 (Protista: Ciliophora: Stichotrichia). – *Zootaxa* **1559**: 31–57.
- Parker J.G. (1981): Ciliated protozoa of the polluted Tees Estuary. – *Estuar. Coastal Shelf Sci.* **12**: 337–340.
- Patterson D.J., Larsen J., Corliss J.O. (1989): The ecology of heterotrophic flagellates and ciliates living in marine sediments. – *Progr. Protistol.* **3**: 185–277.
- Pereyaslawzewa S. (1886): Protozoaires de la mer Noire. – *Zap. novoross. Obshch. Estest.* **10**: 79–114, 3 plates (in Russian).
- Perez-Romero P., Villalobo E., Torres A. (2001): Different stop codon usage in two pseudohypotrich ciliates. – *FEMS Microbiology Letters* **205**: 259–263.
- Perty M. (1849): Mikroskopische Organismen der Alpen und der italienischen Schweiz. – *Mitt. naturf. Ges. Bern year 1849*: 153–176.
- Petran A. (1967): Cercetari asupra faunei de ciliate psamobionte la plajele dn sudul litoralului romanesc al marii negre. – *Ecol. Mar.* **2**: 169–191.
- Petran A. (1968): Sur l'écologie des ciliés psammobiontes de la Mer Noire (littoral Roumain). – *Revue roum. Biol., Zoologie* **13**: 441–446.
- Petran A. (1971): Sur la faune des ciliés des sédiments sablonneux du littoral Roumain de la Mer Noire. – *Cercetari marine* **1**: 149–166.
- Petz W. (1999): Ciliophora. In: Boltovskoy D. (ed.): *South Atlantic Zooplankton*, pp. 265–319. Backhuys Publishers, Leiden.
- Petz W., Foissner W. (1993): Morphogenesis in some freshwater tintinnids (Ciliophora, Oligotrichida). – *Europ. J. Protistol.* **29**: 106–120.
- Petz W., Foissner W. (1996): Morphology and morphogenesis of *Lamtostylia edaphoni* Berger and Foissner and *Onychodromopsis flexilis* Stokes, two hypotrichs (Protozoa: Ciliophora) from Antarctic soils. – *Acta Protozool.* **35**: 257–280.
- Petz W., Foissner W. (1997a): Ecology and taxonomy of terrestrial ciliates of continental Antarctica. – *J. Euk. Microbiol. Suppl.* **44**: 30A, Abstract 120.

- Petz W., Foissner W. (1997b): Morphology and infraciliature of some soil ciliates (Protozoa, Ciliophora) from continental Antarctica, with notes on the morphogenesis of *Sterkiella histriomuscorum*. – Polar Record **33** (187): 307–326.
- Petz W., Wirmsberger E., Foissner W. (1988): Die Kleintiere in den Fichtenwaldböden des Oberen Mühlviertels. Leicht zu übersehen, aber sehr wichtig. In: Amt der Oberösterreichischen Landesregierung, Abteilung Kultur (ed.): Das Mühlviertel. Natur, Kultur, Leben, pp. 77–88. Wimmer-Druck, Linz.
- Petz W., Song W., Wilbert N. (1995): Taxonomy and ecology of the ciliate fauna (Protozoa, Ciliophora) in the endopagial and pelagial of the Weddell Sea, Antarctica. – Stapfia **40**: 1–223.
- Pfannkuche O., Jelinek H., Hartwig E. (1975): Zur Fauna eines Süßwasserwattes im Elbe-Aestuar. – Arch. Hydrobiol. **76**: 475–498.
- Pinto C. (1925): Protozoários observados no Brasil. – Mems Inst. Oswaldo Cruz **18**: 211–301, Estampas 16–28.
- Prescott D.M. (1994): The DNA of ciliated protozoa. – Microbiol. Rev. **58**: 233–267.
- Prescott D.M. (1998): Invention and mystery in hypotricha DNA. – J. Euk. Microbiol. **45**: 575–581.
- Provasoli L., Balamuth W., Becker E.R., Corliss J.O., Hall R.P., Holz G.G.Jr., Levine N.D., Manwell R.D., Nigrelli R.F., Starr R.C., Trager W. (= The Committee on Cultures of the Society of Protozoology) (1958): A catalogue of laboratory strains of free-living and parasitic Protozoa (with sources from which they may be obtained and directions for their maintenance). – J. Protozool. **5**: 1–38.
- Quennerstedt A. (1867): Bidrag till sveriges infusorie-fauna. II. – Acta Univ. lund. **4**: 1–48, Plates I, II (in Swedish).
- Quennerstedt A. (1869): Bidrag till sveriges infusorie-fauna. III. – Acta Univ. lund. **6**: 1–35, 1 plate (in Swedish).
- Raffaelli G.R. (1970): La structure fine de *Diophrys scutum* (Dujardin). – Archs Anat. microsc. Morph. exp. **59**: 221–234.
- Raikov I.B. (1960): La faune interstitielle des infusoires du littoral sableux de la baie Dolniye Zelentzy, Mourmanie. – Trav. Inst. Biol., Mourmansk **2**: 172–185 (in Russian; reference from Hartwig 1973b).
- Raikov I.B. (1962): Les ciliés mésopsammiques du littoral de la Mer Blanche (U.R.S.S.) avec une description de quelques espèces nouvelles ou peu connues. – Cah. Biol. mar. **3**: 325–361.
- Raikov I.B. (1963): Ciliates of the mesopsammon of the Ussuri Gulf (Japan Sea). – Zool. Zh. **42**: 1753–1767 (in Russian with English summary).
- Raikov I.B. (1969): The macronucleus of ciliates. In: Chen T.-T. (ed.): Research in Protozoology Volume 3, pp. 1–128. Pergamon Press, Oxford, London, Edinburgh, New York, Toronto, Sydney, Paris, Braunschweig.
- Raikov I.B. (1982): The protozoan nucleus. Morphology and evolution. In: Alfert M., Beermann W., Goldstein L., Porter K.R., Sitté P. (eds.): Cell biology monographs, Volume 9. Springer, Wien, New York. I–XV, 474 pp.
- Raikov I.B. (1996): Nuclei of ciliates. In: Hausmann K., Bradbury P.C. (eds.): Ciliates. Cells as organisms, pp. 221–242. Gustav Fischer Verlag, Stuttgart, Jena, Lübeck, Ulm.
- Raikov I.B., Kovaleva V.G. (1968): Complements to the fauna of psammobiotic ciliates of the Japan Sea (Posjet gulf). – Acta Protozool. **6**: 309–333.
- Rawlinson N.G., Gates M.A. (1985): The excystment process in the ciliate *Euplotes muscicola*: an integrated light and scanning electron microscopic study. – J. Protozool. **32**: 729–735.
- Rawlinson N.G., Gates M.A. (1986): A morphological classification of encysting species of *Euplotes* (Ciliophora: Nassophorea: Euplotida). – Trans. am. microsc. Soc. **105**: 301–310.
- Rees J.van (1884): Protozën der Oosterschelde (Protozoaires de l'escaut de l'est). – Tijdschr. ned. dierk. Vereen. **Supplement Deel I** (years 1883–1884): 592–673, Plaat XVI (in Dutch with French translation).
- Repak A.J., Anderson O.R. (1990): The fine structure of the encysting salt marsh heterotrich ciliate *Fabrea salina*. – J. Morph. **205**: 335–341.

- Riess F. (1840): Beiträge zur Fauna der Infusorien, mit dem beigegeführten Ehrenberg'schen Systeme. – Dissertation Universität Wien, 40 pp.
- Robinson B.S., Bamforth S.S., Dobson P.J. (2002): Density and diversity of protozoa in some arid Australian soils. – *J. Euk. Microbiol.* **49**: 449–453.
- Ruini J. (1938): Notizen über Ciliaten aus konzentrierten Salzwässern. – *Zoöl. Meded., Leiden* **20**: 243–256.
- Ruiz M.S., Anadón R. (1987): Fines structure of the contractile vacuole complex of *Opercularia coarctata* Claparede & Lachmann (peritrich ciliate). – *Arch. Protistenk.* **133**: 269–275.
- Ruthmann A., Kuck A. (1985): Formation of the cyst wall of the ciliate *Copoda steinii*. – *J. Protozool.* **32**: 677–682.
- Ryan P.G., Watkins B.P., Lewis Smith R.I., Dastych H., Eicker A., Foissner W., Heatwole H., Miller W.R., Thompson G. (1989): Biological survey of Robertsollen, western Dronning Maud Land: area description and preliminary species lists. – *S. Afr. T. Antarkt.* **19**: 10–20.
- Saburova M.A., Azovskii A.I., Polikarpov I.G. (2004): A comparative analysis of the vertical migrations of interstitial ciliates in different types of sediments. – *Oceanology* **44**: 535–547.
- Santangelo G., Lucchesi P. (1995): Spatial distribution pattern of ciliated protozoa in a Mediterranean interstitial environment. – *Aquat. microb. Ecol.* **9**: 47–54.
- Schewiakoff W. (1892): Ueber die geographische Verbreitung der Süßwasser-Protozoen. – *Verh. naturh.-med. Ver. Heidelb. (N. S.)* **4**: 544–567.
- Schewiakoff W. (1893): Über die geographische Verbreitung der Süßwasser-Protozoen. – *Zap. imp. Akad. Nauk, 7<sup>e</sup> Série* **41**: 1–201, Tafeln I–IV.
- Schiewer U. (1994): Vorkommen und Bedeutung der Protozoen in der Darß-Zingster Boddenkette. – *Rostocker Meeresbiologische Beiträge* **2**: 77–83.
- Schmidt S.L., Bernhard D., Foissner W., Berger H., Schlegel M. (2004): Phylogenetische Analyse der Spirotrichea (Ciliophora) mittels Gensequenzen der small subunit (ssu) rDNA. – *Berichte des naturwissenschaftlich-medizinischen Vereins in Innsbruck Supplementum* **15**: 63, Abstract.
- Schmidt S.L., Bernhard D., Schlegel M., Foissner W. (2007): Phylogeny of the Stichotrichia (Ciliophora; Spirotrichea) reconstructed with nuclear small subunit rRNA gene sequences: discrepancies and accordance with morphological data. – *J. Euk. Microbiol.* **54**: 201–209.
- Schröder O. (1906): Beiträge zur Kenntnis von *Campanella umbellaria* L. sp. (*Epistylis flavicans* + *grandis* Ehrbg.). – *Arch. Protistenk.* **7**: 75–105, Tafeln I, II.
- Schwarz M.V.J., Frenzel P. (2003): Population dynamics and ecology of ciliates (Protozoa, Ciliophora) in an anoxic rice field soil. – *Biol. Fertil. Soils* **38**: 245–252.
- Seyffert W. (2003) (ed.): *Lehrbuch der Genetik*. Spektrum Akademischer Verlag, Heidelberg, Berlin. I–XXVIII, 1230 pp.
- Shao C., Song W., Warren A., AL-Rasheid K.A.S., Yi Z., Gong J. (2006): Morphogenesis of the marine ciliate, *Pseudoamphisiella alveolata* (Kahl, 1932) Song & Warren, 2000 (Ciliophora, Stichotrichia, Urostylelida) during binary fission. – *J. Euk. Microbiol.* **53**: 388–396.
- Shao C., Song W., Yi Z., Gong J., Li J., Lin X. (2007): Morphogenesis of the marine spirotrichous ciliate, *Trachelostyla pediculiformis* (Cohn, 1866) Borrer, 1972 (Ciliophora, Stichotrichia), with consideration of its phylogenetic position. – *Europ. J. Protistol.* **43**: 255–264.
- Shi X. (1993): How to distinguish the natural groups in hypotrichous ciliates. – *Int. Congr. Protozool.* **9**: 116, Abstract 454.
- Shi X.-L. (1999): Systematic revision of the order Hypotrichida I. Protohypotrichina and Stichotrichina (Ciliophora). – *Acta Zootax. sinica* **24**: 241–264 (in Chinese with English summary).
- Shi X.-L. (2000): Systematic revision of the order Hypotrichida III. Sporadotrichina and Euplotina (Ciliophora). – *Acta Zootax. sinica* **25**: 9–25 (in Chinese with English summary).
- Shi X., Song W., Shi, X. (1999): Systematic revision of the hypotrichous ciliates. In: Song W. (ed.). *Progress in Protozoology*, pp. 77–154. Qingdao Ocean University Press, Qingdao (in Chinese).

- Shi X., Hu X., Warren A. (2003): Rediscovery of *Gastrostyla setifera* (Engelmann, 1862) Kent, 1882: morphology and divisional morphogenesis (Ciliophora, Hypotrichida). – *J. nat. Hist. (London)* **37**: 1411–1422.
- Shin M.K. (1994): Systematics of Korean hypotrichs (Ciliophora, Polyhymenophora, Hypotrichida) and molecular evolution of hypotrichs. Dissertation Seoul National University, Seoul. 270 pp.
- Sich H. (1990): Die benthische Ciliatenfauna bei Gabelsflach (Kieler Bucht) und deren Beeinflussung durch Bakterien - Eine Studie über Menge, Biomasse, Produktion, Bakterieningestion und Ultrastruktur von Mikroorganismen. – *Ber. Inst. Meeresk. Christian-Albrechts-Universität Kiel* **Nr. 191**: I–IX, 1–215.
- Šlapeta J., Moreira D., López-García P. (2005): The extent of protist diversity: insights from molecular ecology of freshwater eukaryotes. – *Proc. R. Soc. Series B.* **272**: 2073–2081.
- Small E.B., Lynn D.H. (1985): Phylum Ciliophora Doflein, 1901. In: Lee J.J., Hutner S.H., Bovee E.C. (eds.): *An illustrated guide to the protozoa*, pp. 393–575. Society of Protozoologists, Lawrence.
- Smith J.C. (1904): A preliminary contribution to the protozoan fauna of the Gulf Biologic Station with notes on some rare species. – *Rep. Louisiana Gulf Biologic Station for 1903* **2** (year 1904): 43–55.
- Song W. (1990): A comparative analysis of the morphology and morphogenesis of *Gonostomum strenua* (Engelmann, 1862) (Ciliophora, Hypotrichida) and related species. – *J. Protozool.* **37**: 249–257.
- Song W. (2001): Morphology and morphogenesis of the marine ciliate *Ponturostyla enigmatica* (Dragesco & Dragesco-Kernéis, 1986) Jankowski, 1989 (Ciliophora, Hypotrichida, Oxytrichidae). – *Europ. J. Protistol.* **37**: 181–197.
- Song W., Hu X. (1999): Divisional morphogenesis in the marine ciliate, *Hemigastrostyla enigmatica* (Dragesco & Dragesco-Kernéis, 1986) and redefinition of the genus *Hemigastrostyla* Song & Wilbert, 1997 (Protozoa, Ciliophora). – *Hydrobiologia* **391**: 249–257.
- Song W., Packroff G. (1993): Beitrag zur Morphogenese des marinen Ciliaten *Diophrys scutum* (Dujardin, 1841) (Ciliophora, Hypotrichida). – *Zool. Jb. Anat.* **123**: 85–95.
- Song W., Packroff G. (1997): Taxonomische Untersuchungen an marinen Ciliaten aus China mit Beschreibungen von zwei neuen Arten, *Strombidium globosaneum* nov. spec. and *S. platum* nov. spec. (Protozoa, Ciliophora). – *Arch. Protistenk.* **147** (year 1996/97): 331–360.
- Song W., Warren A. (1996): A redescription of the marine ciliates *Uroleptus retractilis* (Claparède and Lachmann, 1858) comb. n. and *Epiclintes ambiguus* (Müller, 1786) Bütschli, 1889 (Ciliophora, Hypotrichida). – *Acta Protozool.* **35**: 227–234.
- Song W., Wilbert N. (1989): Morphology and infraciliature of *Uroleptooides qingdaoensis* sp. nov. (Ciliophora; Hypotrichida: Amphisiellidae). – *Acta Zootax. sinica* **14**: 390–395 (in Chinese with English summary).
- Song W., Wilbert N. (1994): Morphogenesis of the marine ciliate *Diophrys oligothrix* Borror, 1965 during the cell division (Protozoa, Ciliophora, Hypotrichida). – *Europ. J. Protistol.* **30**: 38–44.
- Song W., Wilbert N. (1997): Morphological investigations on some free living ciliates (Protozoa, Ciliophora) from China Sea with description of a new hypotrichous genus, *Hemigastrostyla* nov. gen. – *Arch. Protistenk.* **148**: 413–444.
- Song W., Wilbert N. (1997a): Morphological studies on some free living ciliates (Ciliophora: Heterotrichida, Hypotrichida) from marine biotopes in Qingdao, China, with descriptions of three new species: *Holosticha warreni* nov. spec., *Tachysoma ovata* nov. spec. and *T. dragescoi* nov. spec. – *Europ. J. Protistol.* **33**: 48–62.
- Song W., Wilbert N. (2000): Ciliates from Antarctic sea ice. – *Polar Biol.* **23**: 212–222.
- Song W., Warren A., Hu X. (1997): Morphology and morphogenesis of *Pseudoamphisiella lacazei* (Maupas, 1883) Song, 1996 with suggestion of establishment of a new family Pseudoamphisiellidae nov. fam. (Ciliophora, Hypotrichida). – *Arch. Protistenk.* **147**: 265–276.
- Song W., Wilbert N., Chen Z., Shi X. (2004): Consideration on the systematic position of *Uronychia* and related euplotids based on data of ontogeny and 18S rRNA gene sequence analyses, with morphogenetic redescription of *Uronychia setigera* Calkins, 1902 (Ciliophora: Euplotida). – *Acta Protozool.* **43**: 313–328.

- Song W., Wilbert N., AL-Rasheid K., Warren A., Shao C., Long H., Yi Z., Li L. (2007): Redescriptions of two marine hypotrichous ciliates, *Diophrys irmgard* and *D. hystrix* (Ciliophora, Euplotida), with a brief revision of the genus *Diophrys*. – J. Euk. Microbiol. **54**: 283–296.
- Šrámek-Hušek R. (1954): Neue und wenig bekannte Ciliaten aus der Tschechoslowakei und ihre Stellung im Saprobiensystem. – Arch. Protistenk. **100**: 246–267.
- Stein F. (1854): Die Infusionsthierie auf ihre Entwicklungsgeschichte untersucht. Engelmann, Leipzig. i–x, 265 pp, Tafeln I–VI.
- Stein, F. (1859): Der Organismus der Infusionsthierie nach eigenen Forschungen in systematischer Reihenfolge bearbeitet. I. Abtheilung. Allgemeiner Theil und Naturgeschichte der hypotrichen Infusionsthierie. Engelmann, Leipzig. I–XII, 206 pp, Tafeln I–XIV.
- Stein F. (1859a): Charakteristik neuer Infusorien-Gattungen. – Lotos **9**: 2–5, 57–60.
- Stein F. (1863): Neue Infusorienformen in der Ostsee. – Amtliche Berichte Deutscher Naturforscher und Aerzte in Karlsbad **37. Versammlung im September 1862**: 161–162, 165–166.
- Stein F. (1864): Über den *Proteus tenax* von O. F. Müller und über die Infusoriengattungen *Distigma* Ehrbg. und *Epiclintes* Stein. – Sber. K. böhm. Ges. Wiss. **year 1864**: 41–46.
- Stein F. (1867): Der Organismus der Infusionsthierie nach eigenen Forschungen in systematischer Reihenfolge bearbeitet. II. Abtheilung. 1) Darstellung der neuesten Forschungsergebnisse über Bau, Fortpflanzung und Entwicklung der Infusionsthierie. 2) Naturgeschichte der heterotrichen Infusorien. Engelmann, Leipzig. I–VIII, 355 pp & Tafeln I–XVI.
- Sterki V. (1878): Beiträge zur Morphologie der Oxytrichinen. – Z. wiss. Zool. **31**: 29–58, Tafel IV.
- Stiller J. (1974): Járólábacsok csillósok - Hypotrichida. – Fauna Hung. **115**: 1–187 (in Hungarian).
- Stiller J. (1974a): Ergänzungen der von Fauré-Fremiet vorgenommenen Neuordnung der hypotrichen Ciliaten. – Annl. hist.-nat. Mus. natn. hung. **66**: 129–133.
- Stokes A.C. (1886): Some new hypotrichous infusoria. – Proc. Am. phil. Soc. **23**: 21–30, 1 plate.
- Stokes A.C. (1887): Some new hypotrichous infusoria from American fresh waters. – Ann. Mag. nat. Hist., Serie 5 **20**: 104–114, Plate III.
- Sudhaus W., Rehfeld K. (1992): Einführung in die Phylogenetik und Systematik. Gustav Fischer Verlag, Stuttgart, Jena, New York. I– XII, 241 pp.
- Sudzuki M. (1979): On the microfauna of the Antarctic region III. Microbiota of the terrestrial interstices. – Mem. nat. Inst. Polar Res., Tokyo **Special Issue 11**: 104–126, Plates 1–10.
- Sui S.-G., Chen Y., Qiu Z.-J., Shi X.-B. (2001): Structure of oral apparatus in *Stylonychia mytilus*. – Acta zool. sin. **47**: 436–441, Plates I, II (in Chinese with English summary).
- Summers F.M. (1935): The division and reorganization of the macronuclei of *Aspidisca lynceus* Müller, *Diophrys appendiculata* Stein, and *Stylonychia pustulata* Ehrbg. – Arch. Protistenk. **85**: 173–208, Plates 5–10.
- Sutherland J.L. (1933): Protozoa from Australian termites. – Quart. J. Microsc. Soc. **76**: 145–173.
- Szabó A. (1999): Protozoological research of alkaline biotopes in Hortobágy National Park (HNP). II. Ciliata fauna of salinized soils. – Acta Biol. Debr. Oecol. Hung. **9**: 243–256.
- Szabó A. (2000): A contribution to the ciliata (Protozoa) fauna of chernozem soils. – Miscellanea Zoologica Hungarica **13**: 11–19.
- Tamás G., Gellért J. (1960): Adatok a Balatoni hidroszammon élővilágának ismeretéhez (Daten zur Erkennung der Lebewelt des Hydrosammons im Balaton). – Acta biol. hung. **27**: 65–73 (in Hungarian with German summary).
- Tirjaková E. (1988): Structures and dynamics of communities of ciliated Protozoa (Ciliophora) in field communities. I. Species composition, group dominance, communities. – Biológia, Bratisl. **43**: 497–503.
- Tirjaková E. (1991): Species composition of soil communities of ciliated protozoa (Ciliophora) in relation to fertilizer type and concentration. – Acta Fac. Rerum nat. Univ. comen., Bratisl., Series Zoologia **34**: 39–43.
- Tirjaková E. (1992): Ciliophora of the riverside zone of the Danube river system. – Acta Fac. Rerum nat. Univ. comen., Bratisl., Series Zoologia **36**: 71–80.

- Tirjaková E. (1997): Nálevníky (Ciliophora) prameniska pod veľ'kym javorníkom (Ciliophora of the spring area below the Veľ'ky Javorník). – *Folia faunistica Slovaca* **2**: 7–22 (in Slovakian with English summary).
- Tirjaková E. (2001): Nálevníky (Ciliophora) niektorých lokalít vo vysokých tatrách (Ciliated protozoa (Ciliophora) of some localities in the High Tatras Mts. (Slovakia)). – *Folia faunistica Slovaca* **6**: 9–17 (in Slovakian with English summary).
- Tirjaková E. (2004): Structure and dynamics of ciliated protozoa communities (Protozoa: Ciliophora) in a mountain stream in the High Tatra Mts, Slovakia. – *Biológia, Bratisl.* **59/Suppl. 15**: 3–9.
- Tirjaková E. (2005): Nálevníky (Ciliophora). In: Majzlan O. (ed.): Fauna devínskej kobyly, pp. 20–23. Edícia APOP, Slovakia (in Slovakian).
- Tirjaková E., Bartosová P. (2004): Prvónálezy nálevníkov (Protozoa: Ciliophora) v odumretej drevnej hmote na území Slovenska (First record of ciliates (Protozoa: Ciliophora) in the decaying wood mass in Slovakia). – *Folia faunistica Slovaca* **9**: 15–23 (in Slovakian with English summary).
- Tirjaková E., Sloukal E. (2004): Structure and dynamics of ciliated protozoa (Protozoa: Ciliophora) in an acidified spring area of the Malé Karpaty Mts. – *Biológia, Bratisl.* **59/Suppl. 15**: 11–21.
- Tirjaková E., Vdacny P. (2004): Prvé nálezy druhov *Holostichides chardezi* a *Australocirrus octonucleatus* (Ciliophora: Hypotrichia) na území Slovenska (First records of two species of ciliates: *Holostichides chardezi* and *Australocirrus octonucleatus* (Ciliophora: Hypotrichia) in Slovakia). – *Folia faunistica Slovaca* **4**: 25–30 (in Slovakian with English summary).
- Tucolesco J. (1962): Écodynamique des infusoires du littoral Roumain de la mer Noire et des bassins salés para-marins. – *Annl. Sci. nat. (Zool.)*, 12<sup>e</sup> série **3**: 785–845.
- Tuculescu I. (1965): Biodinamica lacului Techirghiol. Biocenozozele si geneza namolului. Academiie Republicii Socialiste România, Bucarest. 525 pp.
- Tuffrau M. (1964): Quelques variants techniques del'imprégnation des ciliés par le protéinate d'argent. – *Arch. Zool. exp. gén.* **104**: 186–190.
- Tuffrau M. (1967): Perfectionnements et pratique de la technique d'imprégnation au protargol des infusoires ciliés. – *Protistologica* **3**: 91–98.
- Tuffrau M. (1974): Un nouvel Euplotidae: *Cytharoides balechi* n. g., n. sp., cilié hypotriche de l'antarctique. – *Protistologica* **10**: 311–317.
- Tuffrau M. (1979): Une nouvelle famille d'hypotriches, Kahliellidae n. fam., et ses consequences dans la repartition des Stichotrichina. – *Trans. Am. microsc. Soc.* **98**: 521–528.
- Tuffrau M. (1987): Proposition d'une classification nouvelle de l'ordre Hypotrichida (Protozoa, Ciliophora), fondée sur quelques données récentes. – *Annl. Sci. nat. (Zool.)*, 13<sup>e</sup> Série **8** (years 1986–1987): 111–117.
- Tuffrau M., Fleury A. (1994): Classe des Hypotriches Stein, 1859. – *Traite de Zoologie* **2**: 83–151.
- Vacelet E. (1960): Note preliminaire sur la faune infusorienne des “sables a amphioxus” de la baie de Marseille. – *Recl Trav. Stn mar. Endoume* **20**: 53–57.
- Vacelet E. (1961a): Les cilies de la microfaune des “sables mal cibres” des environs de Marseille. – *Recl Trav. Stn mar. Endoume* **22**: 13–19.
- Vacelet E. (1961b): La faune infusorienne des “sables à amphioxus” des environs de Marseille. – *Bull. Inst. océanogr. Monaco* **58** (N° 1202): 1–12.
- Verhoeven R. (2002): Ciliates in coastal dune soils of different stages of development. – *Europ. J. Soil Biol.* **38**: 187–191.
- Voss H.J. (1989): Vergleichende morphogenetische Untersuchungen von 13 Arten der Gattung *Euplotes* (Ciliophora, Hypotrichida). – *Arch. Protistenk.* **137**: 331–344.
- Voß H.-J. (1992): Morphogenesis in *Amphisiella australis* Blatterer and Foissner, 1988 (Ciliophora, Hypotrichida). – *Europ. J. Protistol.* **28**: 405–414.
- Voß H.-J. (1997): Morphology and morphogenesis of *Parentocirrus hortualis* nov. gen., nov. spec.: a new genus within the redefined family Kahliellidae sensu Eigner 1995 (Ciliophora, Hypotrichida). – *Europ. J. Protistol.* **33**: 30–47.

- Wägele J.W. (2001): Grundlagen der phylogenetischen Systematik. Pfeil, München. 320 pp.
- Walker G.K., Mauget T.K. (1980): Encystment and excystment in hypotrich ciliates II. *Diophrys scutum* and remarks on comparative features. – *Protistologica* **16**: 525–531.
- Walker G.K., Edwards C.A., Suchard S.J. (1989): Encystment in the peritrich ciliate *Telotrochidium heneguyi*. – *Cytobios* **59**: 7–18.
- Wallengren H. (1900): Zur Kenntnis der vergleichenden Morphologie der hypotrichen Infusorien. – *Bih. K. svensk VetenskAkad. Handl.* **26**: 1–31.
- Wang C.C., Nie D. (1932): A survey of the marine protozoa of Amoy. – *Contr. biol. Lab. Sci. Soc. China, Zoological Series* **8**: 285–385.
- Wang C.C., Nie D. (1934): On the marine protozoa of Amoy. – *Proc. Pacif. Sci. Congr.* **5**: 4207–4211.
- Wang C.C., Nie D. (1935): Report on the rare and new species of fresh-water infusoria, part II. – *Sinensia, Shanghai* **6**: 399–524.
- Wang R., Qiu Z., Chen J., Warren A., Song W. (2007): Morphogenesis of the freshwater ciliate *Neokeronopsis spectabilis* (Kahl 1932) Warren et al., 2002, based on a China population (Ciliophora: Urostyletidae). – *J. Euk. Microbiol.* **54**: 184–190.
- Wanick R.C., Da Silva-Neto I.D. (2004): Benthic ciliates from Sepetiba Bay (Rio de Janeiro, Brazil) with the description of *Pseudokeronopsis sepetibensis* n. sp. (Spirotrichea: Urostyletida). – *Zootaxa* **587**: 1–11.
- Warren A., Fyda J., Song W. (2002): The morphology of the poorly-known freshwater urostyletid ciliate *Neokeronopsis spectabilis* (Kahl, 1932) nov. gen. nov. comb., (Ciliophora: Urostyletidae), with notes on its morphogenesis. – *Europ. J. Protistol.* **38**: 195–206.
- Wasik A., Mikolajczyk E. (1992): The morphology and ultrastructure of the Antarctic ciliate, *Cymatocylis convallaria* (Tintinnina). – *Acta Protozool.* **31**: 233–239.
- Weisse J.F. (1848): Fünftes Verzeichniss St. Petersburgischer Infusorien, nebst Beschreibung einer neuen Limnias. – *Bull. Acad. Saint-Petersbourg* **6**: 353–364.
- Weisse T. (2006): Biodiversity of freshwater microorganisms - achievements, problems, and perspectives. – *Polish J. Ecol.* **54**: 633–652.
- Wenzel F. (1953): Die Ciliaten der Moosrasen trocken Standorte. – *Arch. Protistenk.* **99**: 70–141.
- Werner, F.C. (1972): Wortelemente lateinisch-griechischer Fachausdrücke in den biologischen Wissenschaften. Suhrkamp, Baden-Baden. 475 pp.
- Westheide W., Rieger R. (1996): Spezielle Zoologie. Erster Teil: Einzeller und wirbellose Tiere. G. Fischer Verlag, Stuttgart Jena New York. 909 pp.
- Wiackowski K. (1981): Small-scale distribution of psammophilic ciliates. – *Acta hydrobiol., Kraków* **23**: 217–226.
- Wicklow B.J. (1981): Evolution within the order Hypotrichida (Ciliophora, Protozoa): ultrastructure and morphogenesis of *Thigmokeronopsis johadai* (n. gen., n. sp.); phylogeny in the Urostylelina (Jankowski, 1979). – *Protistologica* **17**: 331–351.
- Wicklow B.J. (1982a): Cortical morphogenesis during cell division in *Amphisiella marioni* Gourret and Roeser, 1887 and its bearing on hypotrich phylogeny. – *J. Protozool. Suppl.* **29**: 285A, Abstract.
- Wicklow B.J. (1982b): The Discocephalina (n. subord.): ultrastructure, morphogenesis and evolutionary implications of a group of endemic marine interstitial hypotrichs (Ciliophora, Protozoa). – *Protistologica* **18**: 299–330.
- Wicklow B.J. (1983): Morphogenetic and ultrastructural investigation in the Hypotrichida (Ciliophora, Protozoa): implications in ciliate evolution. – *Diss. Abstr. Int.* **43B**: 2135, Abstract.
- Wicklow B.J. (1983a): Ultrastructure and cortical morphogenesis in the euplotine hypotrich *Certesia quadrinuclata* Fabre-Domergue, 1885 (Ciliophora, Protozoa). – *J. Protozool.* **30**: 256–266.
- Wilbert N. (1975): Eine verbesserte Technik der Protargolimprägung für Ciliaten. – *Mikrokosmos* **64**: 171–179.
- Wilbert N. (1995): Benthic ciliates of salt lakes. – *Acta Protozool.* **34**: 271–288.
- Wilbert N., Kahan D. (1986): *Semiplatyophrya foissneri* nov. gen., nov. spec. und *Perisincirra pori* nov. spec., Ciliaten aus einem Salzboden des Sinai. – *Arch. Protistenk.* **131**: 129–138.

- Wilbert N., Song W. (2005): New contributions to the marine benthic ciliates from the Antarctic area, including description of seven new species (Protozoa, Ciliophora). – *J. nat. Hist. (London)* **39**: 935–973.
- Winston J.E. (1999): Describing species. Practical taxonomic procedure for biologists. Columbia University Press, New York.
- Wirnsberger E., Foissner W., Adam H. (1985): Cortical pattern in non-dividers, dividers and reorganizers of an Austrian population of *Paraurostyla weissei* (Ciliophora, Hypotrichida): a comparative morphological and biometrical study. – *Zool. Scr.* **14**: 1–10.
- Wirnsberger-Aescht E., Foissner W., Foissner I. (1989): Morphogenesis and ultrastructure of the soil ciliate *Engelmanniella mobilis* (Ciliophora, Hypotrichida). – *Europ. J. Protistol.* **24**: 354–368.
- Wirnsberger-Aescht E., Foissner W., Foissner I. (1990): Natural and cultured variability of *Engelmanniella mobilis* (Ciliophora, Hypotrichida); with notes on the ultrastructure of its resting cyst. – *Arch. Protistenk.* **138**: 29–49.
- Wise B.N. (1965): The morphogenetic cycle in *Euplotes eurystomus* and its bearing on problems of ciliate morphogenesis. – *J. Protozool.* **12**: 626–648.
- Wright A.-D.G., Lynn D.H. (1997): Maximum ages of ciliate lineages estimated using a small subunit rRNA molecular clock: crown eukaryotes data back to the Paleoproterozoic. – *Arch. Protistenk.* **148**: 329–341.
- Wrzesniowski A. (1877): Beiträge zur Naturgeschichte der Infusorien. – *Z. wiss. Zool.* **29**: 267–323, Tafeln XIX–XXI.
- Wyatt C.E., Pearson T.H. (1982): The Loch Eil project: population characteristics of ciliate protozoans from organically enriched sea-loch sediments. – *J. exp. mar. Biol. Ecol.* **56**: 279–303.
- Xu D., Song W., Lin X., Warren A. (2006): On two marine oligotrich ciliates, *Spirostrombidium agathae* n. sp. and *S. schizostomum* (Kahl, 1932) n. comb. from China, with a key to the identification of seven well-characterized *Spirostrombidium* spp. (Ciliophora: Oligotrichida). – *Acta Protozool.* **45**: 433–442.
- Xu K., Lei Y. (2007): A new spirofilid ciliate from the Yellow Sea: *Metastrongylidium distichum* n. gen., n. sp. (Ciliophora, Stichotrichia). – *Acta Protozool.* **46**: 121–129.
- Xu K., Song W. (1999): Pathogenetic ciliates from mariculture waters. In: Song W. (ed.): *Progress in protozoology*, pp. 211–268. Qingdao Ocean University Press, Qingdao (in Chinese).
- Yang E.J., Choi J.K., Yoo M.H., Cho B.C., Choi D.H. (2005): Temporal variation in the distributions of the benthic heterotrophic protozoa and their grazing impacts on benthic bacteria and microalgae in the Ganghwa Tidal Flat, Korea. – *The Sea. Journal of the Korean Society of Oceanography* **10**: 19–30 (in Korean with English summary).

# Systematic Index

The index contains all names of hypotrichous ciliates mentioned in the book, including vernacular names for example, amphisiellids; food organisms and other ciliates, except for those mentioned in the general section, are not included. The index is two-sided, that is, species appear both with the genus-group name first (for example, *Amphisiella annulata*) and with the species-group name first (*annulata*, *Amphisiella*). Valid (in my judgement) species and genera are in **boldface italics** print. Valid taxa not revised in the present book, invalid taxa, junior homonyms, synonyms, outdated combinations, incorrect spellings, nomina nuda are given in italics. The scientific name of a subgenus, when used with a binomen or trinomen, must be interpolated in parentheses between the genus-group name and the species-group name (ICZN 1999, Article 6.1). In the following index, these parentheses are omitted to simplify electronic sorting. Thus, the name *Holosticha (Amphisiella) annulata* is listed as *Holosticha Amphisiella annulata*. Note that this name is also listed under “*Amphisiella annulata*, *Holosticha*” and “*annulata*, *Holosticha Amphisiella*”. Suprageneric taxa are represented in normal type, valid ones revised in the present review in **boldface**. **Boldface page number** indicates the location of a valid taxon in the systematic section. “T” indicates the location of the table with the morphometric characterisation; “K” marks the page where a taxon is mentioned in a key (note that some taxa are mentioned two times in one key or in two different keys).

- 18-cirri hypotrich(s) xii, 1, 10, 12, 13, 15–17, 21, 25–27, 34, 45, 47, 71, 80, 104, 140, 164, 289, 352, 390, 410, 471, 473, 482, 507, 530, 533, 578, 582, 589, 593, 603, 614
- 18-cirri oxytrichid(s) 44, 104, 120, 454, 469, 675, 681
- abdita***, ***Afroamphisiella*** 4, 165, 371, 372K, 373, 377, 380T
- abdita*, *Lamtostyla* 165, 372, 377
- acrobates*, *Kahlia* 468
- acuta*, *Amphisiella* 89, 351, 352
- acuta***, ***Paramphisiella*** 5, 72, 89, 255, 322, 352K, 352, 356T
- adami*, *Anteholosticha* 39, 639K
- affine*, *Gonostomum* 38, 53, 471, 476, 478
- affine*, *Trachelostyla* 476
- affinis*, *Oxytricha* 512
- Afroamphisiella*** 78K, 166, 323, 353, 359, 371, 564
- Afroamphisiella abdita*** 4, 165, 371, 372K, 373, 377, 380T
- Afroamphisiella multinucleata*** 4, 8, 74, 247, 372K, 372, 380T
- agiliformis*, *Urosomoida* 579, 581, 582
- agilis*, *Uroleptus* 513, 530
- agilis*, *Urosomoida* 587
- Allotricha* 36, 468, 675, 682
- Allotricha antarctica* 615
- Allotricha mollis* 675
- alpestris*, *Anteholosticha* 636K
- Amphisiellidae 79
- Amphisia* 85, 91, 94
- Amphisia marioni* 91
- Amphisia Oxytricha capitata* 91
- Amphisiella* 85
- Amphisiella milnei* 118
- Amphisiella*** 5, 10, 17, 21, 25, 49–52, 66, 75K, 81, 84, 129, 130, 133, 148, 151, 163, 164, 169, 171, 183, 187, 190, 192, 201, 244, 245, 147, 249, 255, 262, 264, 267, 273, 280, 285, 320, 351, 371, 464, 476, 539, 560, 599, 652, 655, 683
- Amphisiella acuta* 89, 351, 352
- Amphisiella annulata*** 1, 4, 5, 7, 8, 15–17, 47, 72, 86T, 90K, 94, 96, 99, 100, 119, 120, 123, 130, 348, 472, 683, 690, 693
- Amphisiella annulata*, *Holosticha* 101
- Amphisiella antarctica* 89, 129
- Amphisiella anulata* 103
- Amphisiella arenicola* 89, 127
- Amphisiella australis* 4, 89, 165, 167K, 169, 173, 179, 226T, 246K, 263
- Amphisiella binucleata* 261, 262, 264, 267
- Amphisiella binucleata binucleata* 262, 264
- Amphisiella binucleata multicirrata* 89, 262, 263, 267
- Amphisiella capitata*** 4, 81, 90K, 91, 103, 108, 109, 116, 117, 119, 123, 128, 187, 190, 201, 683, 686T
- Amphisiella capitata*, *Holosticha* 93
- Amphisiella dorsicirrata* 89
- Amphisiella edaphoni* 324
- Amphisiella elegans* 89, 164, 165, 192
- Amphisiella faurei* 89
- Amphisiella gellerti* 424, 426
- Amphisiella Holosticha annulata* 100
- Amphisiella kihni* 247

- Amphisiella lithophora* 89  
*Amphisiella longiseris* 90, 245, 249  
*Amphisiella magnigranulosa* 47, 90, 245, 273  
*Amphisiella marioni* 82, 84, 91, 93–95, 97, 100, 108, 244, 247, 683, 685, 686T  
*Amphisiella marioni*, *Holosticha* 93  
*Amphisiella milnei* 119  
***Amphisiella milnei*** 4, 5, 8, 90K, 94, 108, 109, 112, 118, 123  
*Amphisiella milnei*, *Holosticha* 89, 119  
*Amphisiella multinucleata* 90, 245, 255  
*Amphisiella namibiensis* 90, 533, 534, 538  
*Amphisiella oblonga* 463  
*Amphisiella oblonga*, *Holosticha* 464  
***Amphisiella oscensis*** 4, 90, 129, 129K, 130, 133, 654K, 655, 657T  
***Amphisiella ovalis*** 4, 89, 91K, 125  
*Amphisiella Oxytricha capitata* 91  
*Amphisiella perisincirra* 200  
*Amphisiella polycirrata* 90, 245, 285  
*Amphisiella procera* 90, 165, 183  
*Amphisiella quadrinucleata* 90, 165, 190  
*Amphisiella raptans* 90, 245, 252, 253  
*Amphisiella terricola* 90, 245, 279  
*Amphisiella Tetrastyla oblonga* 463  
*Amphisiella thiophaga*, *Holosticha* 130  
***Amphisiella turanica*** 4, 89, 91K, 109, 117, 123  
*Amphisiella vitiphila* 187  
*Amphisiella*, *Holosticha* 48, 85, 90  
amphisiellid(s) 1, 3, 5, 7, 10, 12, 13, 15–19, 21, 22, 28, 38, 41, 45, 47, 48, 51–53, 71, 72, 74, 512, 534, 538, 560, 561, 571, 598, 599, 636, 652, 653, 656  
**Amphisiellidae** 1, 12, 18, 24, 46, 48, 49, 71, 79, 476, 534, 560, 561, 563, 651–653, 659  
***Amphisiellides*** xii, 49–51, 78K, 83, 88, 130, 133, 466, 561, 569, 651  
***Amphisiellides atypicus*** 4, 77, 130, 132, 133, 245, 246K, 561, 567T, 569, 652, 653, 654K, 654, 657T, 662  
*Amphisiellides illuvialis* 561, 562, 569, 652–655  
amphisiellids, core 82  
*Amphysiella* 85  
*Amphysiella milnei* 119  
*Ancystropodium* 139  
***annulata*, *Amphisiella*** 1, 4, 5, 7, 8, 15–17, 47, 72, 86T, 90K, 94, 96, 99, 100, 119, 120, 123, 130, 348, 472, 683, 690, 693  
*annulata*, *Amphisiella Holosticha* 100  
*annulata*, *Holosticha* 89, 100  
*annulata*, *Holosticha Amphisiella* 101  
*antarctica*, *Allotricha* 615  
*antarctica*, *Amphisiella* 89, 129  
***antarctica*, *Caudiamphisiella*** 4, 8, 72, 89, 129K, 129, 133, 135T, 151, 437, 655  
*antarctica*, *Urosomoida* 578  
*antecirrata*, *Anteholosticha* 638K  
*Anteholosticha* 77K, 389, 391, 392, 424, 636, 640, 643, 645, 647, 648  
*Anteholosticha adami* 39, 639K  
*Anteholosticha alpestris* 636K  
*Anteholosticha antecirrata* 638K  
*Anteholosticha arenicola* 637K  
*Anteholosticha australis* 637K  
*Anteholosticha azerbaijanica* 638K, 639K  
*Anteholosticha bergeri* 638K, 647–649  
*Anteholosticha brachysticha* 638K, 639K  
*Anteholosticha brevis* 636K  
*Anteholosticha camerounensis* 637K  
*Anteholosticha distyla* 638K  
*Anteholosticha estuarii* 638K  
*Anteholosticha extensa* 637K  
*Anteholosticha fasciola* 639K  
*Anteholosticha gracilis* 638K, 639K  
*Anteholosticha grisea* 639K  
***Anteholosticha heterocirrata*** 4, 77, 339K, 640K, 640  
*Anteholosticha intermedia* 639K  
*Anteholosticha longissima* 639K  
*Anteholosticha macrostoma* 637K  
*Anteholosticha manca* 639K  
*Anteholosticha mancoidea* 638K  
*Anteholosticha monilata* 637K  
*Anteholosticha multistilata* 638K  
*Anteholosticha muscicola* 637K  
*Anteholosticha oculata* 119, 122, 637K  
*Anteholosticha plurinucleata* 639K  
*Anteholosticha pulchra* 28, 638K  
*Anteholosticha randani* 638K, 639K  
*Anteholosticha scutellum* 639K  
*Anteholosticha sigmoidea* 637K  
*Anteholosticha sphagni* 638K  
*Anteholosticha thononensis* 639K  
***Anteholosticha verrucosa*** 4, 393K, 396T, 402, 424, 426, 427, 636, 640K, 647  
*Anteholosticha violacea* 637K, 639K  
*Anteholosticha vuxgracilis* 637K  
*Anteholosticha warreni* 639K  
*Anteholosticha xanthichroma* 637K, 638K  
*anulata*, *Amphisiella* 103  
*Apoamphisiella* 46, 151, 539  
*Apokeronopsis* 47  
***Apourosomoida*** 5, 48, 52, 71, 75K, 78K, 514, 582, 589, 615  
***Apourosomoida halophila*** 4, 10, 18, 21, 75, 515, 516K, 516, 527T, 530, 582

- Apourosomoida natronophila* 4, 52, 515, 516K, **530**  
*Architricha* 36, 675  
*Architricha indica* 675  
*arenicola*, *Amphisiella* 89, 127  
*arenicola*, *Anteholosticha* 637K  
*arenicola*, *Circinella* 391, 392, 467  
*arenicola*, *Erimophrya* 4, 578, 579K, 579, 583T, **586**, 590, 593  
*arenicola*, *Vermioxytricha* 5, 9, 77, 391, 428, 596, 597T, 599, **600**, 607  
*Aspidisca* 25, 28, 30, 31, 33, 43  
Aspidiscidae 24  
aspidiscids 475  
*atypica*, *Uroleptooides* 466, 653–655  
*atypicus*, *Amphisiellides* 4, 77, 130, 132, 133, 245, 246K, 561, 567T, 569, 652, 653, 654K, **654**, 657T, 662  
*atypicus*, *Uroleptooides* 245, 651, 652, 654, 655  
*auricularis*, *Epiclintes* 493  
*auricularis*, *Oxytricha* 467  
*australis*, *Amphisiella* 4, 89, 165, 167K, 169, 173, **179**, 226T, 246K, 263  
*australis*, *Anteholosticha* 637K  
*australis*, *Lamlostyla* 3, 4, 20, 89, 164, 167K, 167, **169**, 179, 180, 182, 183, 187, 201, 212, 226T, 329, 331, 341  
*Australocirrus* 46  
*azerbaijanica*, *Anteholosticha* 638K, 639K  
*Balladyna* 49, 466  
*Balladyna fusiformis* 466  
*Balladyna parvula* 466  
*Balladynella* 49, 466  
Banylsellidae 467  
*Banyulsella* 49, 466  
*Banyulsella viridis* 466  
Banyulsellidae 467  
*bavariensis*, *Gastrostyla* 138  
*bavariensis*, *Gastrostyla Kleinstyla* 140  
*bavariensis*, *Kleinstyla* 140  
*bergeri*, *Anteholosticha* 638K, 647–649  
*bicirratum*, *Tachysoma* 200  
*Biholosticha discocephalus* 472  
*binucleata binucleata*, *Amphisiella* 262, 264  
*binucleata multicirrata*, *Amphisiella* 89, 262, 263, 267  
*binucleata*, *Amphisiella* 261, 262, 264, 267  
*binucleata*, *Amphisiella binucleata* 262, 264  
*binucleata*, *Uroleptooides* 261, 262, 264  
*binucleatus binucleatus*, *Uroleptooides* 4, 9, 226T, 263K, **264**, 267  
*binucleatus multicirratus*, *Uroleptooides* 4, 9, 89, 226T, 251, 264K, 264, **267**  
*binucleatus*, *Uroleptooides* 72, 180, 183, 246K, 249, 253, **261**, 273  
*binucleatus*, *Uroleptooides binucleatus* 4, 9, 226T, 263K, **264**, 267  
*Bistichella* 13, 77K, 289, 514, **532**, 660  
*Bistichella buitkampii* 4, 75, 193, 533, 534K, **535**, 548, 550T, 554, 661  
*Bistichella humicola* 5, 533, 534K, 548, 550T, **556**, 661  
*Bistichella namibiensis* 4, 90, 285, 533, 534K, **538**, 552T, 552, 544  
*Bistichella procera* 4, 533, 534K, 535, **547**, 550T, 554, 555, 661  
*Bistichella terrestris* 4, 533, 534K, 548, 550T, 552, **554**, 557, 661  
*Bistichella terrestris*-group 548  
*borrori*, *Prodiscocephalus* 44  
*brachysticha*, *Anteholosticha* 638K, 639K  
*brachytona*, *Eschaneustyla* 467  
*brandti*, *Cyttarocyclus* 41  
*brevis*, *Anteholosticha* 636K  
*breviseries*, *Orthoamphisiella* 39, 373, 469, 690  
*bryonicolum*, *Trachelostyla* 475  
*bryophila*, *Trachelochaeta* 470  
*buitkampii*, *Bistichella* 4, 75, 193, 533, 534K, **535**, 548, 550T, 554, 661  
*buitkampii*, *Hemisincirra* 5, 74, 388, 390, 393K, **393**, 396T, 407, 408, 428, 437, 599, 640  
*buitkampii*, *Paraurostyla* 533–535, 661  
*buitkampii*, *Perisincirra* 393, 394  
*buitkampii*, *Pseudouroleptus* 535  
*camerounensis*, *Anteholosticha* 637K  
*canadiensis*, *Trachelostyla* 478  
*capitata*, *Amphisia Oxytricha* 91  
*capitata*, *Amphisiella* 4, 81, 90K, **91**, 103, 108, 109, 116, 117, 119, 123, 128, 187, 190, 201, **683**, 686T  
*capitata*, *Amphisiella Oxytricha* 91  
*capitata*, *Holosticha Amphisiella* 93  
*capitata*, *Oxytricha* 89, 91, 93–95  
*caudata*, *Paramphisiella* 5, 18, 245, 351, 352K, 353, 356T, **358**  
*caudata*, *Trachelostyla* 5, 383, 475, 476, 478, 478K, **494**, 501, 502, 511, 689  
*caudata*, *Uroleptooides* 245, 352, 358, 359  
*caudata*, *Urosoma* 631  
*caudatus caudatus*, *Pseudouroleptus* 662, 662K, **663**, 667T, 669, 671  
*caudatus namibiensis*, *Pseudouroleptus* 312, 662, 663K, 667T, **668**  
*caudatus*, *Pseudouroleptus* 4, 77, 289, 293, 308, 469, 533, 554, 653, 658, 660, 661

- caudatus*, *Pseudouroleptus caudatus* 662, 662K, 663, 667T, 669, 671  
*caudatus*, *Uroleptoides* 359  
**Caudiamphisiella** 75K, 84, 90, **129**, 437, 655, 657  
**Caudiamphisiella antarctica** 4, 8, 72, 89, 129K, **129**, 133, 135T, 151, 437, 655  
*Caudiholosticha* 389, 392, 636  
*Caudiholosticha gracilis* 392  
*Caudiholosticha navicularum* 636K  
*Caudiholosticha setifera* 103, 104  
*Caudiholosticha sylvatica* 120  
*Certesias* 25, 28, 31  
Chlamydodontidae 3  
Choreotrichia 24, 38  
Choreotrichida 24  
*cienskowskii*, *Urosoma* 513, 631  
*ciliophorum*, *Trachelostyla* 475  
*Circinella* 50, 82, 370, 389, 391, 392, 432, 467, 599  
*Circinella arenicola* 391, 392, 467  
*Circinella filiformis* 428, 432  
*Circinella vettersi* 428  
*citrina*, *Uroleptopsis* 118, 225  
*Cladotricha* 49, 467, 517  
*Cladotricha koltzowii* 467  
*Cladotricha variabilis* 516, 517, 529  
Cladotrichidae 163, 244  
*Codonella* 25  
colpodid(s) 30, 39, 40  
*Coniculostomum* 36, 46, 49, 467, 673, 675  
core amphisiellids 82  
*corlissi*, *Meseres* 28  
*cornuta*, *Stichochaeta* 477  
*coronata*, *Gastrostyla Holosticha* 137  
*coronata*, *Holosticha* 137, 138, 141, 148  
*coronata*, *Holosticha Keronopsis* 138  
*coronata*, *Keronopsis Holosticha* 137  
*corsica*, *Stichochaeta* 479, 482, 483, 485, 492  
**Cossothigma** 72K, **382**, 472, 478, 481, 510  
**Cossothigma dubium** 4, 74, 153, 380K, **383**, 386, 387, 506  
*crassa*, *Oxytricha* 47  
*crassum*, *Strongylidium* 297  
*cristata*, *Pseudourostyla* 39  
*Cyrtohymena* 44, 46, 206, 259, 261, 536  
*Cyrtohymena tetracirrata* 464  
*Cyrtostrombidium* 38  
*Cytharoides* 28, 31, 43  
*Cyttarocylis brandti* 41  
*decorata*, *Lamtostyla* 4, 8, 164, 165, 166K, 205, 206, **212**, 218, 226T  
*deserticola*, *Urosomoida* 578  
*diademata*, *Holosticha* 90, 130  
*Diophrys* 22, 25, 26, 28–31, 33–44  
*Diophrys hystrix* 41  
*Diophrys scutum* 33, 35, 40  
discocephalid(s) 13, 22, 25, 26, 28–31, 33–44  
Discocephalidae 25  
*Discocephalus* 25, 31  
*discocephalus*, *Biholosticha* 472  
*distichum*, *Metastrongylidium* 148–151, 153, 160T, 161  
*distyla*, *Anteholosticha* 638K  
*dorsicirrata*, *Amphisiella* 89  
*dorsicirrata*, *Gastrostyla* 140  
*dorsicirrata*, *Kleinstyla* 140  
Dorsomarginalia xi, xii, 18, 21, 26, 28, 37, 39–42, 44–47, 80, 83, 163, 206, 321, 390, 472, 515, 534, 560, 561, 563, 566, 577, 578, 596, 598, 599, 614, 615, 617, 618, 651, 653, 658, 689  
Dorsomarginalia, non-oxytrichid 46, 77, 472, 513, 560, 578  
*dubia*, *Gastrostyla* 383  
*dubia*, *Trachelostyla* 382, 383, 478  
**dubium**, **Cossothigma** 4, 74, 153, 380K, **383**, 386, 387, 506  
*dubium*, *Holosticha Parurosoma* 469  
Dysteriidae 3  
*edaphoni*, *Amphisiella* 324  
*edaphoni*, *Lamtostyla* 163, 165, 201, 323–325  
*edaphoni*, *Lamtostylides* 3, 4, 72, 163, 165, 201, 226T, 324K, **324**, 334, 336  
*elegans*, *Amphisiella* 89, 164, 165, 192  
*elegans*, *Lamtostyla* 4, 89, 164, 166K, **192**, 226T  
*elongata*, *Oxytricha* 477  
*elongata*, *Oxytricha Opisthotricha* 477  
*Engelmanniella* 618  
*Engelmanniella mobilis* 42, 607, 618  
*enigmatica*, *Hemigastrostyla* 47, 140, 472, 473, 690  
*enigmatica*, *Mixotricha* 673  
*enigmatica*, *Paraurostyla* 672, 673  
*enigmatica*, *Ponturostyla* 4, 77, 651, **673**, 681T  
*Entodinium* 33, 34  
*Epiclintes* 49, 467  
*Epiclintes auricularis* 493  
**Erimophrya** 71, 78K, 515, 560, **577**, 599, 615, 618, 629  
**Erimophrya arenicola** 4, 578, 579K, 579, 583T, **586**, 590, 593  
**Erimophrya glatzeli** 4, 77, 577, 578, 579K, **579**, 583T, 589  
**Erimophrya quadrinucleata** 4, 516K, 517, 577, 578, 579K, 583T, 592, 593, **595**, 614  
**Erimophrya sylvatica** 4, 578, 579K, 583T, **590**, 595  
*Eschaneustyla* 40, 49

- Eschaneustyla brachytona* 467  
*espeletiae, Fragmocirrus* 193  
*estuarii, Anteholosticha* 638K  
 Euhypotrichina 79  
 Euplota 24, 25  
*Euplotes* 25, 28, 30, 31, 33, 34, 36–40, 43  
 Euplotia 24  
 euplotid(s) 17, 20, 22–26, 28–31, 33–35, 37, 39–44,  
 66, 88, 117, 118, 468, 475  
 Euplotida 24  
 Euplotidae 24  
*Euplotidium* 31  
*extensa, Anteholosticha* 637K  
  
*fasciola, Anteholosticha* 639K  
*faurei, Amphisiella* 89  
*faurei, Marginotricha* 30, 33, 34, 89  
*faurei, Psammocephalus* 33, 89  
*filiformis, Circinella* 428, 432  
*filiformis, Perisincirra* 392  
*flexilis, Onychodromopsis* 468, 615  
*fossicola, Paraurostyla* 539  
*Fragmocirrus espeletiae* 193  
*furcata, Tachysoma* 200  
*fusififormis, Balladyna* 466  
  
 Gastrocirrhidae 24  
*Gastrocirrhus* 31, 138  
*Gastrostyla* 45, 46, 50, 82, 84, 136–141, 151, 383,  
 384, 467, 476, 665, 690  
*Gastrostyla bavariensis* 138  
*Gastrostyla dorsicirrata* 140  
*Gastrostyla dubia* 383  
*Gastrostyla Gastrostyla* 138, 468  
*Gastrostyla Gastrostyla pulchra* 138  
*Gastrostyla Holosticha coronata* 137  
*Gastrostyla Kleinstyla* 138  
*Gastrostyla Kleinstyla bavariensis* 140  
*Gastrostyla minima* 140  
*Gastrostyla muscorum* 139  
*Gastrostyla mystacea* 138  
*Gastrostyla opisthoclada* 140  
*Gastrostyla pulchra* 84, 137, 140, 693  
*Gastrostyla pulchra, Gastrostyla* 138  
*Gastrostyla setifera* 139  
*Gastrostyla Spetastyla* 138, 468  
*Gastrostyla Spetastyla mystacea* 468  
*Gastrostyla steinii* 82, 84, 138–140, 151, 390  
*Gastrostyla stenocephala* 151, 383  
*Gastrostyla Stylonychia pulchra* 137  
*Gastrostyla, Gastrostyla* 138, 468  
 gastrostylid(s) 139, 390  
 Gastrostylidae 139, 140, 390  
*geleii, Trachelostyla* 475  
  
*gellerti gellerti, Hemisincirra* 424, 647  
*gellerti verrucosa, Hemisincirra* 391, 424, 647, 648  
*gellerti, Amphisiella* 424, 426  
*gellerti, Hemisincirra* 388  
***gellerti, Hemisincirra*** 4, 5, 8, 393K, 396T, 418,  
**424**, 437, 647, 648  
*gellerti, Hemisincirra gellerti* 424, 647  
*gellerti, Perisincirra* 424  
*gellerti, Perisincirra* 388, 391, 424, 647  
*gibba, Holosticha* 42  
***glatzeli, Erimophrya*** 4, 77, 577, 578, 579K, **579**,  
 583T, 589  
*Glaucoma* 38  
*Glaucoma scintillans* 53  
***goertzi, Hemiurosoma*** 4, 614, 616T, 618, 619K, **628**  
 Gonostomatidae 471, 472  
*Gonostomum* xii, 16, 40, 45–47, 51, 80, 81, 222,  
 307, 390, 407, 415, 448, 469, 471–473, 475–477,  
 479, 481, 482, 486, 503, 505, 512, 515, 533, 614,  
 615, 617, 618, 623, 625, 631, 691  
*Gonostomum affine* 38, 53, 471, 476, 478  
*Gonostomum namibiense* 690  
*Gonostomum pediculiforme* 479  
*Gonostomum pediculiformis* 479  
*Gonostomum strenuum* 690  
*gracilis, Anteholosticha* 638K, 639K  
*gracilis, Caudiholosticha* 392  
*gracilis, Perisincirra* 388, 392  
*grandis, Onychodromus* 468  
*grandis, Pleurotricha* 675  
*grandis, Urostyla* 41, 42  
***granulifera, Hemiamplesiella*** 4, 8, 289, 290T,  
 294K, 311, **315**, 662  
***granulifera, Lamtostyla*** 4, 8, 10, 16, 48, 72, 164,  
 165, 166K, 205, **206**, 212, 218, 220, 222, 223,  
 226T, 336, 473, 482, 486  
*granulifera, Oxytricha* 482, 615, 689  
***granulifera-group, Lamtostyla*** 71, 78K, 164, 165,  
**205**  
*granuliferum, Strongylidium* 294, 315  
*grelli, Orthoamplesiella* 373  
*grisea, Anteholosticha* 639K  
 group, *Bistichella terrestris*- 548  
**group, Lamtostyla granulifera-** 71, 78K, 164, 165,  
**205**  
**group, Lamtostyla lamottei-** 164, 165, **167**  
**group, Lamtostyla longa-** 71, 78K, 164, 165, **218**  
 group, *Spirotrachelostyla simplex*- 506  
  
*halophila, Apourosomoida* 4, 10, 18, 21, 75, 515,  
 516K, **516**, 527T, 530, 582  
*halophila, Lamtostyla* 165, 323, 336  
*halophila, Parakahiella* 193

- halophilus*, *Lamtostylides* 4, 8, 165, 226T, 323, 324K, **336**, 344, 345, 347, 449, 449K
- Halteria* 24, 33, 37
- halterid(s) 28, 33, 38
- Halteriida 38
- haptorid(s) 30, 39
- hembergeri*, *Spiroamphiella* 4, 17, 72, 140, 148, 149T, **150**, 160T, 384, 506
- Hemiamphiella* 21, 49–51, 77K, 80, 83, 88, 161, 244, **288**, 352, 442, 443, 446, 650, 662, 665, 671
- Hemiamphiella granulifera* 4, 8, 289, 290T, 294K, 311, **315**, 662
- Hemiamphiella qingdaoensis* 310
- Hemiamphiella quadrinucleata* 4, 8, 10, 187, 190, 245, 289, 290T, 294K, **318**, 562, 562K, 563, 564, 567T
- Hemiamphiella terricola* 72, 288, 289, 294K, **294**, 311, 312, 320–322, 352, 442, 562, 660, 690
- Hemiamphiella terricola qingdaoensis* 5, 245, 290T, 295, 296K, **310**
- Hemiamphiella terricola terricola* 5, 8, 289, 290T, 293–295, 296K, **296**, 311
- Hemiamphiella wilberti* 4, 289, 290T, 294K, **311**, 315, 660, 662, 671
- Hemicincirra* sp. 388
- Hemigastrostyla* 141, 151, 384, 472, 473, 691
- Hemigastrostyla enigmatica* 47, 140, 472, 473, 690
- Hemigastrostyla stenocephala* 140, 383, 472
- Hemigastrostyla szaboi* 472
- Hemiscincirra gellerti* 388
- Hemiscincirra* 51, 71, 78K, 83, 139, 200, 201, 323, **387**, 448–450, 458, 512, 578, 579, 598, 599, 607, 614, 618, 633, 634, 636, 643, 645
- Hemiscincirra buitkampii* 5, 74, 388, 390, 393K, **393**, 396T, 407, 408, 428, 437, 599, 643
- Hemiscincirra gellerti* 4, 5, 8, 393K, 396T, 418, **424**, 437, 647, 648
- Hemiscincirra gellerti gellerti* 424, 647
- Hemiscincirra gellerti verrucosa* 391, 424, 647, 648
- Hemiscincirra heterocirrata* 391, 640, 643
- Hemiscincirra inquieta* 5, 8, 390, 391, 393K, 394, 396T, **403**, 437, 473
- Hemiscincirra interrupta* 5, 389, 393K, 396T, 428, **432**, 600, 607
- Hemiscincirra livida* 391, 449, 458
- Hemiscincirra muelleri* 391, 599, 607
- Hemiscincirra namibiensis* 4, 193, 391, 392K, 396T, **418**
- Hemiscincirra octonucleata* 4, 391, 392K, 396T, **421**, 450
- Hemiscincirra polynucleata* 391, 618, 634
- Hemiscincirra pori* 323, 344
- Hemiscincirra quadrinucleata* 4, 193, 245, 391, 392K, 396T, 418, **419**, 449K, 454
- Hemiscincirra rariseta* 5, 389, 391, 393K, 396T, **428**, 599, 600, 607
- Hemiscincirra similis* 633
- Hemiscincirra vermiculare* 435
- Hemiscincirra vermicularis* 5, 389, 391, 392K, 396T, 428, **435**, 599
- Hemiscincirra verrucosa* 647
- Hemiscincirra vettersi* 391
- Hemiscincirra viridis* 450
- Hemiscincirra wenzeli* 5, 8, 391, 393K, 394, 396T, 408, 424, **437**
- Hemiurosoma* 13, 46, 71, 78K, 389, 391, 392, 437, 515, 560, 578, 579, 599, **614**
- Hemiurosoma goertzi* 4, 614, 616T, 618, 619K, **628**
- Hemiurosoma polynucleatum* 4, 619K, 628, **634**
- Hemiurosoma similis* 4, 587, 614, 619K, 622, 631, **633**
- Hemiurosoma terricola* 4, 77, 614, 616T, 618, 619K, **619**
- heterocirrata*, *Anteholosticha* 4, 77, 339K, 640K, **640**
- heterocirrata*, *Hemiscincirra* 391, 640, 643
- heterocirrata*, *Periscincirra* 640
- heterofoissneri*, *Holosticha* 481
- heterotrichid(s) 30
- Histiculus* 46
- holomilnei*, *Holosticha* 119
- Holosticha* 25, 40, 42, 66, 81, 85, 91, 93, 94, 101, 119, 128, 138, 141, 464, 691
- Holosticha Amphiella* 48, 85, 90
- Holosticha Amphiella annulata* 101
- Holosticha Amphiella capitata* 93
- Holosticha Amphiella marioni* 93
- Holosticha Amphiella milnei* 89, 119
- Holosticha Amphiella oblonga* 464
- Holosticha Amphiella thiophaga* 130
- Holosticha annulata* 89, 100
- Holosticha annulata*, *Amphiella* 100
- Holosticha coronata* 137, 138, 141, 148
- Holosticha coronata*, *Gastrostyla* 137
- Holosticha coronata*, *Keronopsis* 137
- Holosticha diademata* 90, 130
- Holosticha gibba* 42
- Holosticha heterofoissneri* 481
- Holosticha holomilnei* 119
- Holosticha Holosticha milnei* 119, 122
- Holosticha Keronopsis coronata* 138
- Holosticha kessleri* 42
- Holosticha milnei* 122
- Holosticha milnei*, *Holosticha* 119, 122
- Holosticha multistilata* 279

- Holosticha obliqua* 103, 104  
*Holosticha Parurosoma dubium* 469  
*Holosticha pullaster* 28  
*Holosticha setifera* 103  
*Holosticha thiophaga* 90  
 holostichid(s) 81, 163, 225  
 Holostichidae 24, 88, 163, 476  
*Holostichides* 469  
**hospes, Mucotrichidium** 4, 74, 75, 153, **441**  
*hospes, Uroleptus* 440, 441  
**humicola, Bistichella** 5, 533, 534K, 548, 550T, **556**, 661  
*humicola, Pseudouroleptus* 557  
*humicola, Uroleptus* 534, 556, 557, 661  
*hyalina, Lamtostyla* 165, 323, 347  
*hyalina, Tachysoma* 324, 347  
*hyalinum, Tachysoma* 348  
**hyalinus, Lamtostylides** 4, 164, 165, 226T, 324K, **347**  
 hypotrich(s) 1, 3, 5, 7, 10, 12, 16–23, 25, 26, 28, 30, 33–36, 38, 41–44, 51, 52, 65, 66, 117, 294, 390, 472, 515, 517, 560, 562, 563, 688  
 hypotrich(s), 18-cirri xii, 1, 10, 12, 13, 15–17, 21, 25–27, 34, 45, 47, 71, 80, 104, 140, 164, 289, 352, 390, 410, 471, 473, 482, 507, 530, 533, 578, 582, 589, 593, 603, 614  
 Hypotricha 1, 12, 17, 21–26, 28–31, 34–38, 40, 41, 43–48, 66, 75, 79, 80, 81, 220, 294, 472, 477, 503, 514, 533, 534, 560, 596, 615, 617, 660, 689, 691  
 Hypotricha, non-dorsomarginalian 46  
 Hypotrichia 24, 28, 44  
 Hypotrichida 25, 79, 384  
*Hypotrichidium* 38  
*hystrix, Diophrys* 41  
*illuvialis, Amphisiellides* 561, 562, 569, 652–655  
**illuvialis, Nudiamphisiella** 4, 562K, 563, 565, 566, 567T, **569**, 576T, 654K, 654–656  
*indica, Architricha* 675  
**inquieta, Hemisincirra** 5, 8, 390, 391, 393K, 394, 396T, **403**, 437, 473  
*inquieta, Perisincirra* 403  
*intermedia, Anteholosticha* 639K  
**interrupta, Hemisincirra** 5, 389, 393K, 396T, 428, **432**, 600, 607  
**interrupta, Nudiamphisiella** 4, 9, 12, 18, 77, 289, 560, 561, 562K, **562**, 567T, 571, 654K  
*interrupta, Paruroleptus* (?) 432  
*interrupta, Perisincirra* 432  
*interrupta, Perisincirra* 388, 391, 432  
**islandica, Lamtostyla** 4, 164, 165, 166K, **196**, 201, 220, 226T, 325, 334,
- kahli, Perisincirra* 393  
*kahli, Uroleptus* 389, 391, 393, 394, 450, 458, 512, 530  
*Kahlia acrobates* 468  
*Kahliella* 49, 468, 472  
 Kahliellidae xii, 88, 244, 660  
*Keronella* 38  
 Keronopsidae 660  
*Keronopsis* 53, 138  
*Keronopsis coronata, Holosticha* 138  
*Keronopsis Holosticha coronata* 137  
*kessleri, Holosticha* 42  
*kihni, Amphisiella* 247  
**kihni, Uroleptoides** 5, 88, 224, 244, 245, 246K, **246**, 294, 352, 372–374  
*Kiitricha* 31  
*kirkeniensis, Lamtostyla* 165, 323, 333  
**kirkeniensis, Lamtostylides** 4, 165, 226T, 324K, 325, **333**, 336  
*kirkensis, Lamtostyla* 334  
*Kleinstyla* 140  
*Kleinstyla bavariensis* 140  
*Kleinstyla bavariensis, Gastrostyla* 140  
*Kleinstyla dorsicirrata* 140  
*Kleinstyla, Gastrostyla* 138  
*koltzowii, Cladotricha* 467  
*Laboea* 38  
*Lacazea* 49, 468  
*Lacazea ovalis* 468  
*lacazei, Pseudoamphisiella* 117  
**lamottei, Lamtostyla** 4, 72, 88, 161, 163–165, 166K, **167**, 171, 187, 190, 201, 220, 245, 512  
**lamottei-group, Lamtostyla** 164, 165, **167**  
**Lamtostyla** 10, 16, 19, 38, 50, 51, 78K, 80, 81, 88, **161**, 245, 246, 273, 322–324, 334, 336, 348, 353, 377, 390, 422, 448, 449, 512  
*Lamtostyla abdita* 165, 372, 377  
**Lamtostyla australis** 3, 4, 20, 89, 164, 167K, 167, **169**, 179, 180, 182, 183, 187, 201, 212, 226T, 329, 331, 341  
**Lamtostyla decorata** 4, 8, 164, 165, 166K, 205, 206, **212**, 218, 226T  
*Lamtostyla edaphoni* 163, 165, 201, 323–325  
**Lamtostyla elegans** 4, 89, 164, 166K, **192**, 226T  
**Lamtostyla granulifera** 4, 8, 10, 16, 48, 72, 164, 165, 166K, 205, **206**, 212, 218, 220, 222, 223, 226T, 336, 473, 482, 486  
**Lamtostyla granulifera-group** 71, 78K, 164, 165, **205**  
*Lamtostyla halophila* 165, 323, 336  
*Lamtostyla hyalina* 165, 323, 347

- Lamostyla islandica* 4, 164, 165, 166K, **196**, 201, 220, 226T, 325, 334  
*Lamostyla kirkeniensis* 165, 323, 333  
*Lamostyla kirkensis* 334  
*Lamostyla lamottei* 4, 72, 88, 161, 163–165, 166K, **167**, 171, 187, 190, 201, 220, 245, 512  
*Lamostyla lamottei-group* 164, 165, **167**  
*Lamostyla longa* 4, 72, 164, 165, 166K, 206, 212, **218**, 222, 223, 226T  
*Lamostyla longa-group* 71, 78K, 164, 165, **218**  
*Lamostyla perisincirra* 3, 4, 164, 166K, 197, 199, **200**, 220, 226T, 329, 341  
*Lamostyla procera* 1, 4, 90, 164, 167K, **183**, 226T, 273  
*Lamostyla quadrinucleata* 4, 90, 164, 166K, 187, **190**, 193, 226T  
*Lamostyla raptans* 4, 164, 166K, 206, 212, 218, 220, **221**, 226T  
*Lamostyla vitiphila* 4, 164, 166K, **187**, 190, 193, 226T, 245, 320  
*Lamostylides* 78K, 164, 166, 246, **322**, 391, 392, 448, 449, 454  
*Lamostylides edaphoni* 3, 4, 72, 163, 165, 201, 226T, 324K, **324**, 334, 336  
*Lamostylides halophilus* 4, 8, 165, 226T, 323, 324K, **336**, 344, 345, 347, 449, 449K  
*Lamostylides hyalinus* 4, 164, 165, 226T, 324K, **347**  
*Lamostylides kirkeniensis* 4, 165, 226T, 324K, 325, **333**, 336  
*Lamostylides pori* 4, 226T, 324K, **344**  
*lanceolata*, *Oxytricha* 617  
*lanceolata*, *Paragastrostyla* 469, 564  
*lanceolata*, *Periholosticha* 469  
*Laurentia monilata* 467  
*Laurentiella* 46, 653  
*lemnae*, *Stylonychia* 42  
*levis*, *Pseudurostyla* 39  
*Limnostrombidium* 25  
*lithophora*, *Amphisiella* 89  
*lithophora*, *Marginotricha* 89  
*lithophora*, *Psammocephalus* 89  
*livida*, *Hemisincirra* 391, 449, 458  
*livida*, *Terricirra* 5, 9, 448, 449K, **458**, 460T  
*Lohmanniella* 25  
*longa*, *Lamostyla* 4, 72, 164, 165, 166K, 206, 212, **218**, 222, 223, 226T  
*longa*, *Tachysoma* 165, 218  
*longa-group*, *Lamostyla* 71, 78K, 164, 165, **218**  
*longiseries*, *Amphisiella* 90, 245, 249  
*longiseries*, *Uroleptoidea* 4, 9, 90, 193, 226T, 246K, **249**  
*longissima*, *Anteholosticha* 639K  
*macrostoma*, *Anteholosticha* 637K  
*macrostoma*, *Trachelostyla* 475  
*magnigranulosa*, *Amphisiella* 47, 90, 245, 273  
*magnigranulosus*, *Uroleptoidea* 4, 9, 47, 90, 182, 183, 226T, 246K, 249, 255, 263, 267, 272, **273**, 390  
*manca*, *Anteholosticha* 639K  
*mancoidea*, *Anteholosticha* 638K  
*Maregastrostyla* 21, 36, 40, 72K, 84, 90, 130, **136**, 151, 468, 690, 693  
*Maregastrostyla pulchra* 4, 9, 36, 72, **137**, 147T, 153, 693  
*Marginotricha* 25  
*Marginotricha faurei* 30, 33, 34, 89  
*Marginotricha lithophora* 89  
*marioni*, *Amphisia* 91  
*marioni*, *Amphisiella* 82, 84, 91, 93–95, 97, 100, 108, 244, 247, 683, 685, 686T  
*marioni*, *Holosticha Amphisiella* 93  
*matsusakai*, *Terricirra* 4, 9, 193, 420, 448, 449, 449K, 450, **454**, 458, 460T  
*Meseres* 24, 33, 37, 39, 40  
*Meseres corlissi* 28  
*Metastrongylidium* 148–151  
*Metastrongylidium distichum* 148–151, 153, 160T, 161  
*milenei*, *Amphisiella* 119  
*milnei*, *Amphisiella* 118  
*milnei*, *Amphisiella* 4, 5, 8, 90K, 94, 108, 109, 112, **118**, 123  
*milnei*, *Amphisiella* 119  
*milnei*, *Holosticha* 122  
*milnei*, *Holosticha Amphisiella* 89, 119  
*milnei*, *Holosticha Holosticha* 119, 122  
*minima*, *Gastrostyla* 140  
*minima*, *Spetastyla mystacea* 140  
*minima*, *Urosomoida* 515, 516K, 517  
*Mitra radiosa* 469, 512  
*Mixotricha* 672, 673, 675  
*Mixotricha enigmatica* 673  
*mobilis*, *Engelmanniella* 42, 607, 618  
*mollis*, *Allotricha* 675  
*monilata*, *Anteholosticha* 637K  
*monilata*, *Laurentia* 467  
*monostyla*, *Urosomoida* 578  
*mucicola*, *Strongylidium* 445  
*Mucotrichidium hospes* 4, 74, 75, 153, **441**  
*Mucotrichidium* 50, 77K, **440**  
*muelleri*, *Hemisincirra* 391, 599, 607  
*muelleri*, *Vermioxytricha* 5, 428, 597T, 598–600, 600K, 605, **607**

- multicirrata*, *Amphisiella binucleata* 89, 262, 263, 267  
*multicirratu*s, *Uroleptoides binucleatus* 4, 9, 89, 226T, 251, 264K, 264, 267  
*multinucleata*, *Afroamphisiella* 4, 8, 74, 247, 372K, 372, 380T  
*multinucleata*, *Amphisiella* 90, 245, 255  
*multinucleatus*, *Uroleptoides* 5, 9, 77, 90, 226T, 246K, 251, 253, 255, 272  
*multistilata*, *Anteholosticha* 638K  
*multistilata*, *Holosticha* 279  
*muscolica*, *Anteholosticha* 637K  
*muscorum*, *Gastrostyla* 139  
*muscorum*, *Strongyliidium* 289, 294, 296, 297, 311  
*musculus*, *Trichoda* 443  
*mutusakai*, *Terircirra* 448  
*mystacea minima*, *Spetastyla* 140  
*mystacea mystacea*, *Spetastyla* 140  
*mystacea*, *Gastrostyla* 138  
*mystacea*, *Gastrostyla Spetastyla* 468  
*mystacea*, *Oxytricha* 140  
*mystacea*, *Spetastyla* 140  
*mystacea*, *Spetastyla mystacea* 140  
*mytilus*, *Stylonychia* 482  
  
*namibiense*, *Gonostomum* 690  
*namibiensis*, *Amphisiella* 90, 533, 534, 538  
*namibiensis*, *Bistichella* 4, 90, 285, 533, 534K, 538, 552T, 552, 554  
*namibiensis*, *Hemisincirra* 4, 193, 391, 392K, 396T, 418  
*namibiensis*, *Pseudouroleptus caudatus* 312, 662, 663K, 667T, 668  
*natronophila*, *Apourosomoida* 4, 52, 515, 516K, 530  
*natronophilus*, *Uroleptus* 515, 530, 531  
*navicularum*, *Caudiholosticha* 636K  
*Neokeronopsis* xi, 25, 44, 46, 47, 65, 82, 359, 560, 661  
*Neokeronopsis spectabilis* 18  
non-dorsomarginalian *Hypotricha* 46  
non-oxytrichid *Dorsomarginalia* 46, 77, 472, 513, 560, 578  
non-stylonychine *Oxytrichidae* 46  
*Notohymena* 46  
*nova*, *Pseudourostyla* 636  
*Nudiamphisiella* 21, 46, 78K, 83, 289, 320, 321, 560, 561, 653, 655  
*Nudiamphisiella illuvialis* 4, 562K, 563, 565, 566, 567T, 569, 576T, 654K, 654–656  
*Nudiamphisiella interrupta* 4, 9, 12, 18, 77, 289, 560, 561, 562K, 562, 567T, 571, 654K  
*obliqua*, *Holosticha* 103, 104  
  
*oblonga*, *Amphisiella* 463  
*oblonga*, *Amphisiella Tetrastyla* 463  
*oblonga*, *Holosticha Amphisiella* 464  
*oblonga*, *Tetrastyla* 4, 74, 75, 463  
*octonucleata*, *Hemisincirra* 4, 391, 392K, 396T, 421, 450  
*octonucleata*, *Perisincirra* 421  
*octonucleata*, *Urosoma* 614  
*oculata*, *Anteholosticha* 119, 122, 637K  
*oligotrich(s)* 23–26, 28–31, 33, 34, 36–44, 117  
*Oligotricha* 24, 25, 37  
*Oligotrichea* 24, 38, 41  
*Oligotrichia* 24, 37, 38  
*Oligotrichida* 24, 41  
*Onychodromopsis* 49, 468  
*Onychodromopsis flexilis* 468, 615  
*Onychodromus* 23, 46, 49, 468, 653  
*Onychodromus grandis* 468  
*Onychodromus quadricornutus* 468  
*Ophrydium* 447  
*opisthoclada*, *Gastrostyla* 140  
*Opisthotricha elongata*, *Oxytricha* 477  
*Orthoamphisiella* 50, 51, 82, 190, 244, 371, 377, 468  
*Orthoamphisiella breviseries* 39, 373, 469, 690  
*Orthoamphisiella grelli* 373  
*Orthoamphisiella stramenticola* 373, 468  
*Orthoamphisiellidae* 467, 469  
*oscensis*, *Amphisiella* 4, 90, 129, 129K, 130, 133, 654K, 655, 657T  
*ovalis*, *Amphisiella* 4, 89, 91K, 125  
*ovalis*, *Lacazea* 468  
*Oxytricha* 23, 25, 33, 34, 38, 42, 43, 46, 81, 104, 112, 220, 473, 577, 596, 615, 659, 689–691  
*Oxytricha affinis* 512  
*Oxytricha auricularis* 467  
*Oxytricha capitata* 89, 91, 93–95  
*Oxytricha capitata*, *Amphisia* 91  
*Oxytricha capitata*, *Amphisiella* 91  
*Oxytricha crassa* 47  
*Oxytricha elongata* 477  
*Oxytricha granulifera* 482, 615, 689  
*Oxytricha lanceolata* 617  
*Oxytricha mystacea* 140  
*Oxytricha Opisthotricha elongata* 477  
*Oxytricha setigera* 466  
*Oxytricha stenocephala* 141  
*Oxytrichia* 24  
oxytrichid(s) xii, 12, 18–21, 24, 26, 28, 34, 37, 39, 40, 42, 45, 51, 65, 71, 81–83, 88, 138, 140, 141, 151, 163, 206, 289, 293, 390, 466, 468–470, 472, 473, 475, 477, 513, 515, 533, 539, 560, 561, 578,

- 582, 596, 615, 651–653, 659–661, 665, 673, 675, 682, 688, 689
- oxytrichid(s), 18-cirri 44, 104, 120, 454, 469, 675, 681
- Oxytrichidae xi, 12, 18, 24–26, 29, 42, 44–47, 77, 81, 82, 88, 140, 162, 163, 293, 384, 447, 448, 467, 470, 473, 475, 514, 515, 533, 560, 577, 578, 596, 598, 615, **651**, 653, 660, 672
- Oxytrichidae, non-stylonychine 46
- Oxytrichidea 24
- Oxytrichinae 44, 47
- oxytrichines 45
- Parabirojimia similis* 690
- Paragastrostyla* 50, 51, 392, 469, 564
- Paragastrostyla lanceolata* 469, 564
- Paragonostomum* 473
- Paraholosticha* 53
- Parakahliella* 36, 46
- Parakahliella halophila* 193
- Parakahliellidae 653
- Paramecium* 33, 34, 38
- Paramphisiella*** 49–51, 78K, 88, 244, 294, 322, **351**, 352
- Paramphisiella acuta*** 5, 72, 89, 255, 322, 352K, **352**, 356T
- Paramphisiella caudata*** 5, 18, 245, 351, 352K, 353, 356T, **358**
- Parastylonychia* 139
- Paraurostyla* 33, 34, 43, 46, 49, 469, 535, 560, 673
- Paraurostyla buitkampii* 533–535, 661
- Paraurostyla enigmatica* 672, 673
- Paraurostyla fossicola* 539
- Paraurostyla viridis* 618
- Paraurostyla weissei* 42, 535, 673
- Parentocirrus* 539
- Paruroleptus* (?) *interrupta* 432
- Paruroleptus* 225
- Parurosoma* 49, 469
- Parurosoma dubium*, *Holosticha* 469
- parvula*, *Balladyna* 466
- Pattersoniella* xi, 45–47, 653, 691
- paucicirrata*, *Periholosticha* 407
- pediculariformis*, *Trachelostyla* 480
- pediculiforme*, *Gonostomum* 479
- pediculiforme*, *Stichochoaeta* 493
- pediculiforme*, *Trachelostyla* 480
- pediculiformis*, *Gonostomum* 479
- pediculiformis*, *Stichochoaeta* 4, 470, 474, 475, 477, 478, 500
- pediculiformis*, *Trachelostyla*** 5, 12, 18, 26, 75, 383, 384, 386, 387, 390, 476, 478K, **478**, 494–498, 500, 501T, 502, 505, 683, **687**
- pediculiformis*, *Trachelostyla Stichochoaeta* 479, 480
- pediculiformis*, *Traxhelostyla* 475
- Pelagohalteria* 24
- Pelagotrichidium* 38
- pellionellum*, *Tachysoma* 5, 120
- pendiculiformis*, *Trachelostyla* 480
- Periholosticha* 51, 391, 392, 407, 469, 643
- Periholosticha lanceolata* 469
- Periholosticha paucicirrata* 407
- Perilemmaphora*** 25, 26, **37**, 38, 40, 42
- Perisincira gellerti* 424
- Perisincira interrupta* 432
- Perisincira viridis* 450
- Perisincirra* 200, 387–391, 394, 424, 432, 450, 634
- Perisincirra buitkampii* 393, 394
- Perisincirra filiformis* 392
- Perisincirra gellerti* 388, 391, 424, 647
- Perisincirra gracilis* 388, 392
- Perisincirra heterocirrata* 640
- Perisincirra inquieta* 403
- Perisincirra interrupta* 388, 391, 432
- Perisincirra kahli* 393
- Perisincirra octonucleata* 421
- Perisincirra pori* 324, 344
- Perisincirra quadrinucleata* 419
- Perisincirra similis* 392, 618, 631
- Perisincirra vermiculare* 435
- Perisincirra viridis* 392, 447–450, 512
- perisincirra*, *Amphisiella* 200
- perisincirra*, *Lamtostyla*** 3, 4, 164, 166K, 197, 199, **200**, 220, 226T, 329, 341
- perisincirra*, *Tachysoma* 165, 196, 200
- peritrich(s) 28
- Petalotricha* 41
- Phacodinium* 29, 31, 34, 36, 40
- piscis*, *Uroleptus* 447, 617
- Pleurotricha* 46, 675
- Pleurotricha grandis* 675
- Pleurotricha setifera* 139
- plurinucleata*, *Anteholosticha* 639K
- polycirrata*, *Amphisiella* 90, 245, 285
- polycirratus*, *Uroleptoidea*** 4, 90, 226T, 246K, **285**, 539
- Polyhymenophora 24
- polynucleata*, *Hemisincirra* 391, 618, 634
- polynucleata*, *Urosoma* 634
- polynucleatum*, *Hemiurosoma*** 4, 619K, 628, **634**
- Ponturostyla*** 21, 36, 78K, 651, **672**
- Ponturostyla enigmatica*** 4, 77, 651, **673**, 681T
- pori*, *Hemisincirra* 323, 344
- pori*, *Lamtostylides*** 4, 226T, 324K, **344**
- pori*, *Perisincirra* 324, 344
- procera*, *Amphisiella* 90, 165, 183

- procera*, *Bistichella* 4, 533, 534K, 535, **547**, 550T, 554, 555, 661
- procera*, *Lamtoystyla* 1, 4, 90, 164, 167K, **183**, 226T, 273
- procerus*, *Pseudouroleptus* 534, 535, 547, 661
- Prodiscocephalus* 25, 31
- Prodiscocephalus borrori* 44
- Protocruzia* 23
- Protogastrostyla* 137, 138, **690**
- Protogastrostyla pulchra* 683, 691T, **693**
- Psammocephalus faurei* 33, 89
- Psammocephalus lithophora* 89
- Psammomitra* 49–51, 469, 471, 473, 512
- Psammomitra retractilis* 469
- Pseudoamphisiella* 118
- Pseudoamphisiella lacazei* 117
- pseudocrassum*, *Strongylidium* 83
- pseudokeronopsid(s)* 40
- Pseudokeronopsidae* 477
- Pseudokeronopsinae* 30, 65
- Pseudokeronopsis* 47
- Pseudokeronopsis qingdaoensis* 47
- Pseudokeronopsis rubra* 279
- Pseudorurostyla levis* 39
- Pseudostrombidium* 38
- Pseudouroleptus* xii, 18, 21, 46, 49, 50, 77K, 79, 83, 151, 289, 293, 307, 312, 469, 470, 533, 535, 548, 554, 557, 651, 653, **658**
- Pseudouroleptus buitkampii* 535
- Pseudouroleptus caudatus* 4, 77, 289, 293, 308, 469, 533, 554, 653, 658, 660, 661
- Pseudouroleptus caudatus caudatus* 662, 662K, **663**, 667T, 669, 671
- Pseudouroleptus caudatus namibiensis* 312, 662, 663K, 667T, **668**
- Pseudouroleptus humicola* 557
- Pseudouroleptus procerus* 534, 535, 547, 661
- Pseudouroleptus terrestris* 534, 554, 557, 661
- Pseudourostyla* 36, 675, 682
- Pseudourostyla cristata* 39
- Pseudourostyla nova* 636
- Psilotricha succisa* 117
- pulchra*, *Anteholosticha* 28, 638K
- pulchra*, *Gastrostyla* 84, 137, 140, 693
- pulchra*, *Gastrostyla Gastrostyla* 138
- pulchra*, *Gastrostyla Stylonychia* 137
- pulchra*, *Maregastrostyla* 4, 9, 36, 72, **137**, 147T, 153, 693
- pulchra*, *Protogastrostyla* 683, 691T, **693**
- pulchra*, *Stilonichia* 137, 690
- pullaster*, *Holosticha* 28
- pyriformis*, *Tetrahymina* 53
- qingdaoensis*, *Hemiamphisiella* 310
- qingdaoensis*, *Hemiamphisiella terricola* 5, 245, 290T, 295, 296K, **310**
- qingdaoensis*, *Pseudokeronopsis* 47
- qingdaoensis*, *Uroleptooides* 245, 294, 295, 308
- quadricornutus*, *Onychodromus* 468
- quadrinucleata*, *Amphisiella* 90, 165, 190
- quadrinucleata*, *Erimophrya* 4, 516K, 517, 577, 578, 579K, 583T, 592, 593, **595**, 614
- quadrinucleata*, *Hemiamphisiella* 4, 8, 10, 187, 190, 245, 289, 290T, 294K, **318**, 562, 562K, 563, 564, 567T
- quadrinucleata*, *Hemisincirra* 4, 193, 245, 391, 392K, 396T, 418, **419**, 449K, 454
- quadrinucleata*, *Lamtoystyla* 4, 90, 164, 166K, 187, **190**, 193, 226T
- quadrinucleata*, *Perisincirra* 419
- quadrinucleata*, *Uroleptooides* 245, 318
- quadrinucleatus*, *Uroleptooides* 294, 318, 320
- radiosa*, *Mitra* 469, 512
- randani*, *Anteholosticha* 638K, 639K
- raptans*, *Amphisiella* 90, 245, 252, 253
- raptans*, *Lamtoystyla* 4, 164, 166K, 206, 212, 218, 220, **221**, 226T
- raptans*, *Tachysoma* 165, 221
- raptans*, *Uroleptooides* 4, 90, 246K, 249, **252**, 255
- rariseta*, *Hemisincirra* 5, 389, 391, 393K, 396T, **428**, 599, 600, 607
- retractilis*, *Psammomitra* 469
- Retroextendia 40
- Rigidocortex* 46
- Rigidothrix* 1, 29, 38, 65
- Rimostrombidium* 39
- Rootlephorida 89
- rostrata*, *Trachelostyla* 4, 385, 387, 476, 478, 478K, 481, 483, **498**
- rubra*, *Pseudokeronopsis* 279
- Rubrioxytricha* 46
- schiffmanni*, *Wallackia* 470
- scintillans*, *Glaucoma* 53
- scutellum*, *Anteholosticha* 639K
- scutum*, *Diophrys* 33, 35, 40
- setifera*, *Caudiholosticha* 103, 104
- setifera*, *Gastrostyla* 139
- setifera*, *Holosticha* 103
- setifera*, *Pleurotricha* 139
- setigera*, *Oxytricha* 466
- sigmoidea*, *Anteholosticha* 637K
- similis*, *Hemisincirra* 633
- similis*, *Hemiurosoma* 4, 587, 614, 619K, 622, 631, **633**
- similis*, *Parabirojimia* 690

- similis*, *Perisincirra* 392, 618, 631  
*similis*, *Urosoma* 587, 631  
*simplex* *Trachelostyla* 509  
**simplex**, *Spirotrachelostyla* 4, 384, 503K, 506, **509**  
*simplex*, *Stichotricha* 503, 509  
*simplex*, *Strichotricha* 510  
*simplex*, *Stychotricha* 510  
*simplex*-group, *Spirotrachelostyla* 506  
 sp., *Hemicinsirra* 388  
 sp./spec., *Trachelostyla* 4, 382, 383K, 384, **386**, 475, 477, 480, 481, 485, 489, 501T  
 spathidiids 36  
*Spathidium* 33, 34  
 spec./sp., *Uroleptus* 42, 394, 395, 661  
*spectabilis*, *Neokeronopsis* 18  
*Spetastyla* 140  
*Spetastyla mystacea* 140  
*Spetastyla mystacea minima* 140  
*Spetastyla mystacea mystacea* 140  
*Spetastyla mystacea*, *Gastrostyla* 468  
*Spetastyla*, *Gastrostyla* 138, 468  
*sphagni*, *Anteholosticha* 638K  
*spiralis*, *Spirotrachelostyla* 5, 75, 503K, **503**  
*spiralis*, *Trachelostyla* 478, 502–504  
*Spiretella* 38  
***Spiroamphisiella*** 21, 72K, 78K, 84, 90, **148**, 149T, 445, 472, 473, 478, 503, 510  
***Spiroamphisiella hembergeri*** 4, 17, 72, 140, 148, 149T, **150**, 160T, 384, 506  
 Spirofilidae 148, 440, 442  
***Spirotrachelostyla*** 384, 473, 473K, 478, **502**  
***Spirotrachelostyla simplex*** 4, 384, 503K, 506, **509**  
*Spirotrachelostyla simplex*-group 506  
***Spirotrachelostyla spiralis*** 5, 75, 503K, **503**  
***Spirotrachelostyla tani*** 4, 384, 502, 503K, 504, **506**, 507T, 510, 689  
*Spirotricha* 23–25, 37, 42  
 Spirotrichea 24  
*spirotrichoides*, *Trachelostyla* 475  
 spirotrichs xi, 3, 24, 30, 33, 35, 38, 39, 42  
 Sporadotrichina 473  
*Steinia* 46, 603  
*steinii*, *Gastrostyla* 82, 84, 138–140, 151, 390  
*stenocephala*, *Gastrostyla* 151, 383  
*stenocephala*, *Hemigastrostyla* 140, 383, 472  
*stenocephala*, *Oxytricha* 141  
*Sterkiella* 25, 46  
*Stichochaeta* 476, 477, 480, 481  
*Stichochaeta cornuta* 477  
*Stichochaeta corsica* 479, 482, 483, 485, 492  
*Stichochaeta pediculiforme* 493  
*Stichochaeta pediculiformis* 4, 470, 474, 475, 477, 478, 500  
*Stichochaeta pediculiformis*, *Trachelostyla* 479, 480  
*Stichotricha* 384, 472, 476, 477, 481, 501, 503, 509, 510  
*Stichotricha simplex* 503, 509  
 Stichotrichia 24, 38, 391, 691  
 Stichotrichida 24  
 stichotrichines 619  
*Stilonichia pulchra* 137, 690  
*stramenticola*, *Orthoamphisiella* 373, 468  
*stramenticola*, *Territricha* 470  
*strenuom*, *Gonostomum* 690  
*Strichotricha simplex* 510  
 strobiliidiids 117  
*Strobilidium* 25, 30  
*Strobilidium gyrans* 30  
*Strobilidium velox* 41  
 Strombidiida 38  
 strombidiids 117  
*Strombidium* 25, 38  
 strongylidids 83  
<sup>2</sup>*Strongylidium* 83, 149, 297, 311, 315, 445, 503  
<sup>3</sup>*Strongylidium crassum* 297  
*Strongylidium granuliferum* 294, 315  
*Strongylidium mucicola* 445  
*Strongylidium muscorum* 289, 294, 296, 297, 311  
*Strongylidium pseudocrassum* 83  
*Strongylidium wilberti* 294, 311  
*Stychotricha simplex* 510  
*Stylonychia* 17, 23, 25, 33, 34, 38, 43, 46, 137, 489  
*Stylonychia lemnae* 42  
*Stylonychia mytilus* 482  
*Stylonychia pulchra*, *Gastrostyla* 137  
 Stylonychinae 1, 29–31, 44, 46, 47, 65, 82, 138–140, 467, 468, 653  
 stylonychine(s) 7, 30, 38, 44, 45, 65, 138, 139, 390, 653, 675  
*Styloxophrya* 46, 468  
*succisa*, *Psilotricha* 117  
 suctorian(s) 51  
*sylvatica*, *Caudiholosticha* 120  
***sylvatica*, *Erimophrya*** 4, 578, 579K, 583T, **590**, 595  
*szaboi*, *Hemigastrostyla* 472  
*Tachysoma* 201, 220, 222, 348  
*Tachysoma bicirratum* 200  
*Tachysoma furcata* 200  
*Tachysoma hyalina* 324, 347  
*Tachysoma hyalinum* 348  
*Tachysoma longa* 165, 218  
*Tachysoma pellionellum* 5, 120  
*Tachysoma perisincirra* 165, 196, 200  
*Tachysoma raptans* 165, 221

- tani*, *Spirotrachelostyla* 4, 384, 502, 503K, 504, 506, 507T, 510, 689  
*tani*, *Trachelostyla* 478, 503, 506  
*Tericirra mutsusakai* 448  
*terrestris*, *Pseudouroleptus* 534, 554, 557, 661  
*terrestris*, *Bistichella* 4, 533, 534K, 548, 550T, 552, 554, 557, 661  
*terrestris*-group, *Bistichella* 548  
*Terricirra* 51, 77K, 164, 166, 323, 339, 389, 391, 392, 418, 420, 421, 447, 512  
*Terricirra livida* 5, 9, 448, 449K, 458, 460T  
*Terricirra matsusakai* 4, 9, 193, 420, 448, 449, 449K, 450, 454, 458, 460T  
*Terricirra viridis* 4, 9, 74, 420, 422, 448, 449K, 450, 458, 460T  
*terricola qingdaoensis*, *Hemiamphisiella* 5, 245, 290T, 295, 296K, 310  
*terricola terricola*, *Hemiamphisiella* 5, 8, 289, 290T, 293–295, 296K, 296, 311  
*terricola*, *Amphisiella* 90, 245, 279  
*terricola*, *Hemiamphisiella* 72, 288, 289, 294K, 294, 311, 312, 320–322, 352, 442, 562, 660, 690  
*terricola*, *Hemiamphisiella terricola* 5, 8, 289, 290T, 293–295, 296K, 296, 311  
*terricola*, *Hemiurosoma* 4, 77, 614, 616T, 618, 619K, 619  
*terricola*, *Uroleptoidea* 4, 90, 225, 226T, 246K, 279, 285  
*Territricha* xi, 26, 46, 50, 470  
*Territricha stramenticola* 470  
*Tetmemena* 46  
*tetracirrata*, *Cyrtohymena* 464  
*Tetrahymena* 33, 34, 38  
*Tetrahymena pyriformis* 53  
*Tetrastyla* 77K, 463  
*Tetrastyla oblonga* 4, 74, 75, 463  
*Tetrastyla oblonga*, *Amphisiella* 463  
*Thigmokeronopsis* 36, 47  
*thiophaga*, *Holosticha* 90  
*thiophaga*, *Holosticha Amphisiella* 130  
*thononensis*, *Anteholosticha* 639K  
*tintinnid*(s) 28, 41  
*Trachelochaeta* 49, 50, 470, 477  
*Trachelochaeta bryophila* 470  
*Trachelostya* 475  
*Trachelostyla* xii, 12, 40, 47, 50, 51, 141, 383, 384, 390, 448, 470–473, 473K, 474, 502, 503, 506, 510, 685, 688  
*Trachelostyla affine* 476  
*Trachelostyla bryonicolum* 475  
*Trachelostyla canadiensis* 478  
*Trachelostyla caudata* 5, 383, 475, 476, 478, 478K, 494, 501, 502, 511, 689  
*Trachelostyla ciliophorum* 475  
*Trachelostyla dubia* 382, 383, 478  
*Trachelostyla geleii* 475  
*Trachelostyla macrostoma* 475  
*Trachelostyla pediculariformis* 480  
*Trachelostyla pediculiforme* 480  
*Trachelostyla pediculiformis* 5, 12, 18, 26, 75, 383, 384, 386, 387, 390, 476, 478K, 478, 494–498, 500, 501T, 502, 505, 683, 687  
*Trachelostyla pendiculiformis* 480  
*Trachelostyla rostrata* 4, 385, 387, 476, 478, 478K, 481, 483, 498  
*Trachelostyla simplex* 509  
*Trachelostyla sp./spec.* 4, 382, 383K, 384, 386, 475, 477, 480, 481, 485, 489, 501T  
*Trachelostyla spiralis* 478, 502–504  
*Trachelostyla spirotrichoides* 475  
*Trachelostyla Stichochaeta pediculiformis* 479, 480  
*Trachelostyla tani* 478, 503, 506  
*trachelostylid*(s) 4, 10, 16, 24, 51, 52, 75, 448, 471  
*Trachelostylidae* xii, 1, 12, 24, 46, 48, 51, 71, 72K, 163, 390, 448, 470, 471  
*Traxhelostyla pediculiformis* 475  
*Trichoda musculus* 443  
*turanica*, *Amphisiella* 4, 89, 91K, 109, 117, 123  
*uroleptid*(s) 26  
*Uroleptoidea* 19, 49–51, 77K, 81, 88, 161, 165, 166, 180, 183, 187, 224, 294, 310, 318, 320, 352–354, 372, 376, 377, 655  
*Uroleptoidea atypica* 466, 653–655  
*Uroleptoidea atypicus* 245, 651, 652, 654, 655  
*Uroleptoidea binucleata* 261, 262, 264  
*Uroleptoidea binucleatus* 72, 180, 183, 246K, 249, 253, 261, 273  
*Uroleptoidea binucleatus binucleatus* 4, 9, 226T, 263K, 264, 267  
*Uroleptoidea binucleatus multicirratus* 4, 9, 89, 226T, 251, 264K, 264, 267  
*Uroleptoidea caudata* 245, 352, 358, 359  
*Uroleptoidea caudatus* 359  
*Uroleptoidea kihni* 5, 88, 224, 244, 245, 246K, 246, 294, 352, 372–374  
*Uroleptoidea longiseris* 4, 9, 90, 193, 226T, 246K, 249  
*Uroleptoidea magnigranulosus* 4, 9, 47, 90, 182, 183, 226T, 246K, 249, 255, 263, 267, 272, 273, 390  
*Uroleptoidea multinucleatus* 5, 9, 77, 90, 226T, 246K, 251, 253, 255, 272

- Uroleptooides polycirratu*s 4, 90, 226T, 246K, **285**, 539  
*Uroleptooides qingdaoensis* 245, 294, 295, 308  
*Uroleptooides quadrinucleata* 245, 318  
*Uroleptooides quadrinucleatus* 294, 318, 320  
*Uroleptooides raptans* 4, 90, 246K, 249, **252**, 255  
*Uroleptooides terricola* 4, 90, 225, 226T, 246K, **279**, 285  
*Uroleptooides vitiphila* 165, 187, 245  
*Uroleptooides vitiphilus* 187  
*Uroleptopsis* 225  
*Uroleptopsis citrina* 118, 225  
*Uroleptus* xi, 25, 42, 44, 46, 47, 65, 82, 224, 394, 531, 659, 691  
*Uroleptus agilis* 513, 530  
*Uroleptus hospes* 440, 441  
*Uroleptus humicola* 534, 556, 557, 661  
*Uroleptus kahli* 389, 391, 393, 394, 450, 458, 512, 530  
*Uroleptus natronophilus* 515, 530, 531  
*Uroleptus piscis* 447, 617  
*Uroleptus spec./sp.* 42, 394, 395, 661  
*Uronychia* 25, 28, 31, 33, 36, 41, 43  
*Urosoma* 13, 41, 45, 46, 51, 78K, 389, 437, 439, 471, 473, 513–515, 560, 564, 599, 614, 615, 618, 618K, 619, 623, 625, 630, 633, 634  
*Urosoma caudata* 631  
*Urosoma cienkowskii* 513, 631  
*Urosoma octonucleata* 614  
*Urosoma polynucleata* 634  
*Urosoma similis* 587, 631  
*Urosomoida* 45, 46, 178, 471, 473, 513–515, 530, 531, 566, 577, 589, 596, 599, 603, 615, 619, 634  
*Urosomoida agiliiformis* 579, 581, 582  
*Urosomoida agilis* 587  
*Urosomoida antarctica* 578  
*Urosomoida deserticola* 578  
*Urosomoida minima* 515, 516K, 517  
*Urosomoida monostyla* 578  
*Urosomoides* 51, 513  
*Urospinula* 1, 29  
*Urostyla* 25, 618, 672  
*Urostyla grandis* 41, 42  
*Urostyla viridis* 618  
*Urostyla weissei* 469  
*Urostylida* 24  
*Urostylidae* 24, 81  
*Urostylina* 24  
*urostyloid(s)* xii, 3, 12, 13, 15, 17–20, 24–26, 28, 29, 35, 38, 39, 42, 42, 44, 45, 47, 51, 65, 66, 71, 80–82, 88, 94, 119, 163, 390, 391, 445, 469, 470, 476, 477, 560, 564, 569, 615, 636, 648, 661, 675  
*Urostyloidea* xi, 12, 24, 29, 39, 40, 46, 77, 88, 512, **636**, 643  
*variabilis*, *Cladotricha* 516, 517, 529  
*velox*, *Strobilidium* 41  
*vermiculare*, *Hemisincirra* 435  
*vermiculare*, *Perisincirra* 435  
*vermicularis*, *Hemisincirra* 5, 389, 391, 392K, 396T, 428, **435**, 599  
*Vermioxytricha* 46, 71, 78K, 389, 391, 392, 428, 462, 515, 560, 578, 579, **596**, 615, 618  
*Vermioxytricha arenicola* 5, 9, 77, 391, 428, 596, 597T, 599, **600**, 607  
*Vermioxytricha muelleri* 5, 428, 597T, 598–600, 600K, 605, **607**  
*verrucosa*, *Anteholosticha* 4, 393K, 396T, 402, 424, 426, 427, 636, 640K, **647**  
*verrucosa*, *Hemisincirra* 647  
*verrucosa*, *Hemisincirra gellerti* 391, 424, 647, 648  
*vettarsi*, *Circinella* 428  
*vettarsi*, *Hemisincirra* 391  
*violacea*, *Anteholosticha* 637K, 639K  
*viridis*, *Banyulsella* 466  
*viridis*, *Hemisincirra* 450  
*viridis*, *Paraurostyla* 618  
*viridis*, *Perisincirra* 450  
*viridis*, *Perisincirra* 392, 447–450, 512  
*viridis*, *Terricirra* 4, 9, 74, 420, 422, 448, 449K, **450**, 458, 460T  
*viridis*, *Urostyla* 618  
*vitiphila*, *Amphisiella* 187  
*vitiphila*, *Lantostyla* 4, 164, 166K, **187**, 190, 193, 226T, 245, 320  
*vitiphila*, *Uroleptooides* 165, 187, 245  
*vitiphilus*, *Uroleptooides* 187  
*vuxgracilis*, *Anteholosticha* 637K  
*Wallackia* 46, 50, 470, 472, 473  
*Wallackia schiffmanni* 470  
*warreni*, *Anteholosticha* 639K  
*weissei*, *Paraurostyla* 42, 535, 673  
*weissei*, *Urostyla* 469  
*wenzeli*, *Hemisincirra* 5, 8, 391, 393K, 394, 396T, 408, 424, **437**  
*wilberti*, *Hemiamphisiella* 4, 289, 290T, 294K, **311**, 315, 660, 662, 671  
*wilberti*, *Strongylidium* 294, 311  
*xanthichroma*, *Anteholosticha* 637K, 638K

# Table Index

Table 1. . . . .	page 4
Table 2. . . . .	page 8
Table 3. . . . .	page 24
Table 3a. . . . .	page 34
Table 4. . . . .	page 49
Table 5. . . . .	page 49
Table 6. . . . .	page 49
Table 7. . . . .	page 49
Table 8. . . . .	page 49
Table 9. . . . .	page 50
Table 10. . . . .	page 50
Table 11. . . . .	page 50
Table 12. . . . .	page 50
Table 13. . . . .	page 51
Table 14. . . . .	page 51
Table 15. . . . .	page 86
Table 16. . . . .	page 135
Table 17. . . . .	page 147
Table 18. . . . .	page 149
Table 19. . . . .	page 160
Table 20. . . . .	page 171
Table 21. . . . .	page 171
Table 22. . . . .	page 226
Table 23. . . . .	page 290
Table 24. . . . .	page 356
Table 25. . . . .	page 380
Table 26. . . . .	page 396
Table 27. . . . .	page 404
Table 28. . . . .	page 440
Table 29. . . . .	page 460
Table 30. . . . .	page 501
Table 31. . . . .	page 507
Table 32. . . . .	page 527
Table 33. . . . .	page 550
Table 34. . . . .	page 552
Table 35. . . . .	page 567
Table 36. . . . .	page 567
Table 37. . . . .	page 576
Table 38. . . . .	page 583
Table 39. . . . .	page 597
Table 40. . . . .	page 616
Table 41. . . . .	page 657
Table 42. . . . .	page 667
Table 43. . . . .	page 681
Table 44. . . . .	page 686
Table 45. . . . .	page 691